

12

EUROPEAN PATENT APPLICATION

21 Application number: 83112017.5

22 Date of filing: 30.11.83

51 Int. Cl.³: **C 07 C 103/56, C 07 C 103/76;**
C 07 C 103/50, C 07 C 87/50,
C 07 D 295/00, C 07 D 227/00,
C 07 D 233/60, C 07 D 243/08,
A 61 K 31/16, A 61 K 31/195,
A 61 K 31/33

30 Priority: 30.11.82 JP 208678/82

43 Date of publication of application: 13.06.84
Bulletin 84/24

64 Designated Contracting States: AT BE CH DE FR GB IT
LI NL SE

71 Applicant: Eisai Co., Ltd., 6-10, Kolshikawa 4-chome
Bunkyo-ku, Tokyo 112 (JP)

72 Inventor: Yamatsu, Isao, 3605-669, Kashiwada
Ushiku-machi, Inashiki-gun Ibaragi (JP)
Inventor: Suzuki, Takeshi, 56-107, Sakaecho 1 chome
Ushikumachi, Inashiki-gun Ibaragi (JP)
Inventor: Abe, Shinya, 40-2, Shiroyama Kudzakimachi,
Inashiki-gun Ibaragi (JP)
Inventor: Nakamoto, Kouji, 712-91, Ohaza-Nakanuki,
Tsuchiura-shi Ibaragi (JP)
Inventor: Kajiwara, Akiharu, 25-25, Ninomiya 3 chome
Yatabemachi, Tsukuba-gun Ibaragi (JP)
Inventor: Fujimori, Tohru, 9-9, Toukoudai 2 chome
Toyosatomachi, Tsukuba-gun Ibaragi (JP)
Inventor: Harada, Koukichi, 19-14, Kasuga 4 chome
Yatabe-machi, Tsukuba-gun Ibaragi (JP)
Inventor: Kitamura, Shinichi, 34-7-504, Shimura 1 chome,
Itabashi-ku Tokyo (JP)

74 Representative: Hansen, Bernd, Dr.rer.nat. et al,
Hoffmann, Eitle & Partner Patentanwälte
Arabellastrasse 4, D-8000 München 81 (DE)


54 Polyprenyl compound, process for the production thereof and drug containing the same.

57 A novel polyprenyl compound such as a polyprenyl carb-
oxylic acid amide is disclosed. It has antithrombic and anti-
platelet aggregation activity.

EP 0 110 397 A2

Polyprenyl Compound, Process for the
Production thereof and Drug containing
the same


This invention relates to a polyprenyl compound having excellent medicinal activity. More particularly, the invention relates to a polyprenyl compound of the formula (I):



wherein A, B, Y and Z are each hydrogen, or the pair (1) A and B and/or the pair (2) Y and Z together represent a direct valence bond between the carbon atoms to which they are attached, thereby forming a double bond therebetween; W is a group of -COR or a group of X; and n is zero or an integer of 1 to 4 when W is the group of -COR; n is an integer of 1 to 3 when W is the group of X.

R in the formula is selected from :


(1) a group of the formula



0110397


wherein R^1 is hydrogen or lower alkyl and m is an integer of from 1 to 5;

(2) a group of the formula




wherein k and l are the same or different and each is an integer of from 1 to 5;

(3) a group of the formula




15 wherein R^2 is hydrogen, lower alkyl or aryl, preferably alkyl or aryl;

(4) a group of the formula



wherein p is an integer of from 0 to 5 and R^3 and R^4 are each hydrogen or lower alkyl, preferably lower alkyl;

(5) a group of the formula




wherein q is an integer of from 1 to 5, R^5 , R^6 and R^7

0110397


are each hydrogen or lower alkyl, preferably lower alkyl, and X is a halogen;

(6) a group of the formula




wherein r is 2 or 3 and R⁸ is lower alkyl;

10 (7) a group of the formula




wherein s is 2 or 3, R⁹ and R¹⁰ are each lower alkyl and X is a halogen;

(8) a group of the formula.




wherein D is a group of the formula $\text{---(CH}_2\text{)}_t\text{OH}$, in which
25 t is an integer of from 0 to 5, a group of the formula



wherein u is an integer of from 0 to 5 and R¹¹ and R¹²


0110397

are each hydrogen or lower alkyl, or a group of the formula




wherein v is an integer of from 0 to 5, R^{13} , R^{14} and R^{15} are each lower alkyl and X is a halogen;

(9) a group of the formula




wherein R^{16} is hydrogen or lower alkyl and w is an integer of from 1 to 5;

(10) a group of the formula



25 wherein R^{17} is hydrogen or lower alkyl and x is an integer of from 0 to 5, preferably from 1 to 5; and


(11) a group of the formula




wherein R^{18} is hydrogen or lower alkyl and y is an integer of 1 to 5,

0110397

X is selected from
(1) a group of the formula




wherein R²² is a group of the formula




wherein R²³ is hydrogen or lower alkyl and m is an
integer of from 1 to 5;
a group of the formula

0110397




5 wherein k and l are the same or different and each is an integer of from 1 to 5;

a group of the formula



15 wherein p is an integer of from 0 to 5 and R^{24} and R^{25} are each hydrogen or lower alkyl;

and a group of the formula




wherein q and i are each an integer of 1 to 5 and R^{26} , R^{27} , R^{28} and R^{29} are each a lower alkyl, and R^{21} is hydrogen, a lower alkyl or a halogen atom,

25

30


0110397

(2) a group of the formula




5 wherein K and L are both hydrogen or represent a direct valence bond the carbon atoms to which they are attached,


(3) a group of the formula




(4) a group of the formula



20 (5) a group of the formula




(6) a group of the formula



wherein a is zero or an integer of 1 to 5, and R^{30} is a lower alkyl,

0110397


(7) a group of the formula



5

wherein b is zero or an integer of 1 to 5 and R³¹ is a lower alkyl,


(8) a group of the formula



10

wherein c is zero or an integer of 1 to 5,


(9) a group of the formula



15


wherein d is zero or an integer of 1 to 5 and R³² and R³³ are each a lower alkyl,

(10) a group of the formula




25

(11) a group of the formula




wherein R³⁴ is a lower alkyl,

(12) a group of the formula




30

(13) a group of the formula




(14) a group of the formula.




wherein e and f are each an integer of 1 to 5 and R^{35} ,
 10 R^{36} , R^{37} and R^{38} are each hydrogen or a lower alkyl,

(15) a group of the formula



15 wherein g is an integer of 1 to 5 and R^{39} and R^{40} are
 each hydrogen or a lower alkyl, and

(15) a group of the formula



wherein h is an integer of 1 to 5 and R^{41} and R^{42} are
 each hydrogen or a lower alkyl,
 or a pharmaceutically acceptable salt thereof.

25

30

The compound of the invention is called as the first compound group when W in the formula (I) is the group of -COR and as the second compound group when W is the group of X.

5 The invention also relates to a process for the preparation of the compounds of the formula (I) and pharmacologically acceptable salts thereof, and a pharmaceutical composition containing the formula (I) compound.

10 The term "lower alkyl group" as used in the definition of R^1 through R^{18} and R^{21} through R^{42} in the formula (I) means both straight-chain and branched alkyl groups having 1 to 6 carbon atoms, such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, 15 1-methylpropyl, tert-butyl, n-pentyl, 1-ethylpropyl, isoamyl, and n-hexyl. The term "halogen" as used herein means chlorine, bromine, iodine and fluorine. The compounds of the present invention in which (1) the pair A and B and/or (2) the pair Y and Z together form a 20 double bond between the associated adjacent carbon atoms can exist in the form of various stereoisomers, and these stereoisomers are also included within the scope of the present invention.

The compounds (I) of the present invention may form 25 salts depending on the identity of the substituent W. In appropriate cases, the compounds of the present invention can be easily reacted with a pharmacologically acceptable organic or inorganic acid to form acid addition salts. Examples of such inorganic acids are 30 hydrochloric acid, hydrobromic acid, hydriodic acid and sulfuric acid. Examples of such organic acids are maleic acid, fumaric acid, succinic acid, acetic acid, malonic acid, citric acid and benzoic acid.

Examples of typical compounds of the first compound group according to the invention are listed below.

- 5 N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-ethanolamine,
N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-propanolamine,
N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-butyl alcoholamine,
10 N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-amyl alcoholamine,
N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-hexyl alcoholamine,
N-methyl-N-(3,7,11,15-tetramethyl-2,6,10,14-
15 hexadecatetraenoyl)-ethanolamine,
N-methyl-N-(3,7,11,15-tetramethyl-2,6,10,14-
hexadecatetraenoyl)-propanolamine,
N-ethyl-N-(3,7,11,15-tetramethyl-2,6,10,14-
hexadecatetraenoyl)-ethanolamine,
20 N-ethyl-N-(3,7,11,15-tetramethyl-2,6,10,14-
hexadecatetraenoyl)-propanolamine,
N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-diethanolamine,
N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
25 tetraenoyl)-dipropanolamine,
N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-N-8-hydroxyethyl-propanolamine,
N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-glycine,
30 N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-glycine ethyl ester,
N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-glycine propyl ester,

N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
 tetraenoyl)-glycine allyl ester,
 3-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
 tetraenoylamino)-1-ethylpiperidine,
 5 2-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
 tetraenoylaminomethyl)-1-ethylpyrrolidine,
 N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
 tetraenoyl)-ethylenediamine,
 10 N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
 tetraenoyl)-ethylenediamine hydrochloride,
 N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
 tetraenoyl)-N',N'-dimethylethylenediamine,
 N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
 tetraenoyl)-N',N'-diethylethylenediamine,
 15 N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
 tetraenoyl)-N'-methyl-N'-ethylethylenediamine,
 N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
 tetraenoyl)-N',N'-dimethyl-1,3-diaminopropane,
 20 N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
 tetraenoyl)-N',N'-diethyl-1,3-diaminopropane,
 N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
 tetraenoyl)-N'-methyl-N'-ethyl-1,3-diamino-
 propane,
 25 N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
 tetraenoyl)-N',N',N'-trimethylethylenediamine
 chloride,
 N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
 tetraenoyl)-N',N',N'-trimethylethylenediamine
 iodide,
 30 N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
 tetraenoyl)-N',N',N'-triethylethylenediamine
 chloride,

0110397

- N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-N',N',N'-triethylethylenediamine
iodide,
5 1-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-4-methylpiperazine,
1-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-4-ethylpiperazine,
1-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-4-propylpiperazine,
10 1-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-4-methyl-hexahydro-1,4-diazepine,
1-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-4-ethyl-hexahydro-1,4-diazepine,
15 1-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-4-propyl-hexahydro-1,4-diazepine,
1-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-4,4-dimethylpiperazine chloride,
1-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-4,4-diethylpiperazine chloride,
20 N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-3-hydroxypiperidine,
N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-2-hydroxymethylpiperidine,
N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-3-(dimethylamino)-piperidine,
25 N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-3-(diethylamino)-piperidine,
N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-2-(dimethylaminomethyl)-piperidine,
30 N-(3,7,11,15-tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-2-(diethylaminomethyl)-piperidine,
N-(3,7,11,15-tetramethyl-hexadecanoyl)-
ethanolamine,

0110397

N-(3,7,11,15-tetramethyl-hexadecanoyl)-
propanolamine,
N-(3,7,11,15-tetramethyl-hexadecanoyl)-
diethanolamine,
5 N-(3,7,11,15-tetramethyl-hexadecanoyl)-glycine,
N-(3,7,11,15-tetramethyl-hexadecanoyl)-
ethylenediamine,
N-(3,7,11,15-tetramethyl-hexadecanoyl)-
ethylenediamine hydrochloride,
10 N-(3,7,11,15-tetramethyl-hexadecanoyl)-N'-N'-
dimethylethylenediamine,
N-(3,7,11,15-tetramethyl-hexadecanoyl)-N',N',N'-
trimethylethylenediamine chloride,
N-(3,7,11,15-tetramethyl-hexadecanoyl)-3-
15 hydroxypiperidine,
N-(3,7,11,15-tetramethyl-hexadecanoyl)-2-
hydroxymethylpiperidine,
N-(3,7,11,15-tetramethyl-hexadecanoyl)-3-
(dimethylamino)-piperidine,
20 N-(3,7,11,15-tetramethyl-hexadecanoyl)-2-
(dimethylaminomethyl)-piperidine,
1-(3,7,11,15-tetramethyl-hexadecanoyl)-4-
methylpiperazine,
1-(3,7,11,15-tetramethyl-hexadecanoyl)-4-
25 methyl-hexahydro-1,4-diazepine,
N-methyl-N-(3,7,11,15-tetramethyl-hexadecanoyl)-
ethanolamine,
3-(3,7,11,15-tetramethyl-hexadecanoylamino)-
1-ethylpiperidine,
30 2-(3,7,11,15-tetramethyl-hexadecanoylamino-
methyl)-1-ethylpyrrolidine,
N-(3,7,11,15-tetramethyl-2-hexadecenoyl)-
ethanolamine,

0110397

- N-(3,7,11,15-tetramethyl-2-hexadecenoyl)-
propanolamine,
N-(3,7,11,15-tetramethyl-2-hexadecenoyl)-
diethanolamine,
5 N-(3,7,11,15-tetramethyl-2-hexadecenoyl)-
glycine,
N-(3,7,11,15-tetramethyl-2-hexadecenoyl)-
ethylenediamine,
10 N-(3,7,11,15-tetramethyl-2-hexadecenoyl)-
ethylenediamine hydrochloride,
N-(3,7,11,15-tetramethyl-2-hexadecenoyl)-
N',N'-dimethylethylenediamine,
N-(3,7,11,15-tetramethyl-2-hexadecenoyl)-
N',N',N'-trimethylethylenediamine chloride,
15 N-(3,7,11,15-tetramethyl-2-hexadecenoyl)-
3-hydroxypiperidine,
N-(3,7,11,15-tetramethyl-2-hexadecenoyl)-
2-hydroxymethylpiperidine,
20 N-(3,7,11,15-tetramethyl-2-hexadecenoyl)-
3-(dimethylamino)-piperidine,
N-(3,7,11,15-tetramethyl-2-hexadecenoyl)-
2-(dimethylaminomethyl)-piperidine,
1-(3,7,11,15-tetramethyl-2-hexadecenoyl)-
4-methylpiperidine,
25 1-(3,7,11,15-tetramethyl-2-hexadecenoyl)-
4-methyl-hexahydro-1,4-diazepine,
N-methyl-N-(3,7,11,15-tetramethyl-2-hexade-
cenoyl)-ethanolamine,
3-(3,7,11,15-tetramethyl-2-hexadecenoylamino)-
30 1-ethylpiperidine,
2-(3,7,11,15-tetramethyl-2-hexadecenoylamino-
methyl)-1-ethylpyrrolidine,

0110397

- N-(3,7,11,15-tetramethyl-6,10,14-hexadecatrienoyl)-ethanolamine,
N-(3,7,11,15-tetramethyl-6,10,14-hexadecatrienoyl)-propanolamine,
5 N-(3,7,11,15-tetramethyl-6,10,14-hexadecatrienoyl)-diethanolamine,
N-(3,7,11,15-tetramethyl-6,10,14-hexadecatrienoyl)-glycine,
10 N-(3,7,11,15-tetramethyl-6,10,14-hexadecatrienoyl)-ethylenediamine,
N-(3,7,11,15-tetramethyl-6,10,14-hexadecatrienoyl)-ethylenediamine hydrochloride,
N-(3,7,11,15-tetramethyl-6,10,14-hexadecatrienoyl)-N',N'-dimethylethylenediamine,
15 N-(3,7,11,15-tetramethyl-6,10,14-hexadecatrienoyl)-N',N',N'-trimethylethylenediamine chloride,
N-(3,7,11,15-tetramethyl-6,10,14-hexadecatrienoyl)-3-hydroxypiperidine, -
20 N-(3,7,11,15-tetramethyl-6,10,14-hexadecatrienoyl)-2-hydroxymethylpiperidine,
N-(3,7,11,15-tetramethyl-6,10,14-hexadecatrienoyl)-3-(dimethylamino)-piperidine,
25 N-(3,7,11,15-tetramethyl-6,10,14-hexadecatrienoyl)-2-(dimethylaminomethyl)-piperidine,
1-(3,7,11,15-tetramethyl-6,10,14-hexadecatrienoyl)-4-methylpiperazine,
1-(3,7,11,15-tetramethyl-6,10,14-hexadecatrienoyl)-4-methyl-hexahydro-1,4-diazepine,
30 N-methyl-N-(3,7,11,15-tetramethyl-6,10,14-hexadecatrienoyl)-ethanolamine,
3-(3,7,11,15-tetramethyl-6,10,14-hexadecatrienoylamino)-1-ethylpiperidine,

0110397

- 2-(3,7,11,15-tetramethyl-6,10,14-hexadeca-
trienoylaminomethyl)-1-ethylpyrrolidine,
N-(3,7,11-trimethyl-2,6,10-dodecatrienoyl)-
ethanolamine,
5 N-(3,7,11-trimethyl-2,6,10-dodecatrienoyl)-
propanolamine,
N-(3,7,11-trimethyl-2,6,10-dodecatrienoyl)-
diethanolamine,
N-(3,7,11-trimethyl-2,6,10-dodecatrienoyl)-
10 glycine,
N-(3,7,11-trimethyl-2,6,10-dodecatrienoyl)-
ethylenediamine,
N-(3,7,11-trimethyl-2,6,10-dodecatrienoyl)-
ethylenediamine hydrochloride,
15 N-(3,7,11-trimethyl-2,6,10-dodecatrienoyl)-
N',N'-dimethylethylenediamine,
N-(3,7,11-trimethyl-2,6,10-dodecatrienoyl)-
N',N',N'-trimethylethylenediamine chloride,
N-(3,7,11-trimethyl-2,6,10-dodecatrienoyl)-
20 3-hydroxypiperidine,
N-(3,7,11-trimethyl-2,6,10-dodecatrienoyl)-
2-hydroxypiperidine,
N-(3,7,11-trimethyl-2,6,10-dodecatrienoyl)-
3-dimethylamino-piperidine,
25 N-(3,7,11-trimethyl-2,6,10-dodecatrienoyl)-
2-dimethylaminomethyl-piperidine,
1-(3,7,11-trimethyl-2,6,10-dodecatrienoyl)-
4-methylpiperazine,
1-(3,7,11-trimethyl-2,6,10-dodecatrienoyl)-
30 4-methyl-hexahydro-1,4-diazepine,
N-methyl-N-(3,7,11-trimethyl-2,6,10-dodeca-
trienoyl)-ethanolamine,

0110397

- 3-(3,7,11-trimethyl-2,6,10-dodecatrienoylamino)-
1-ethylpiperidine,
2-(3,7,11-trimethyl-2,6,10-dodecatrienoylamino-
methyl)-1-ethylpyrrolidine,
5 N-(3,7,11,15,19-pentamethyl-2,6,10,14,18-
eicosapentaenoyl)-ethanolamine,
N-(3,7,11,15,19-pentamethyl-2,6,10,14,18-
eicosapentaenoyl)-propanolamine,
10 N-(3,7,11,15,19-pentamethyl-2,6,10,14,18-
eicosapentaenoyl)-diethanolamine,
N-(3,7,11,15,19-pentamethyl-2,6,10,14,18-
eicosapentaenoyl)-glycine,
N-(3,7,11,15,19-pentamethyl-2,6,10,14,18-
eicosapentaenoyl)-ethylenediamine,
15 N-(3,7,11,15,19-pentamethyl-2,6,10,14,18-
eicosapentaenoyl)-ethylenediamine hydrochloride,
N-(3,7,11,15,19-pentamethyl-2,6,10,14,18-
eicosapentaenoyl)-N',N'-dimethylethylenediamine,
20 N-(3,7,11,15,19-pentamethyl-2,6,10,14,18-
eicosapentaenoyl)-N',N',N'-trimethylethylene-
diamine-chloride,
N-(3,7,11,15,19-pentamethyl-2,6,10,14,18-
eicosapentaenoyl)-3-hydroxypiperidine,
25 N-(3,7,11,15,19-pentamethyl-2,6,10,14,18-
eicosapentaenoyl)-2-hydroxymethylpiperidine,
N-(3,7,11,15,19-pentamethyl-2,6,10,14,18-
eicosapentaenoyl)-3-dimethylaminopiperidine,
N-(3,7,11,15,19-pentamethyl-2,6,10,14,18-
eicosapentaenoyl)-2-dimethylaminomethyl-
30 piperidine,
N-(3,7,11,15,19-pentamethyl-2,6,10,14,18-
eicosapentaenoyl)-3-diethylaminopiperidine,

N-(3,7,11,15,19-pentamethyl-2,6,10,14,18-
eicosapentaenoyl)-2-diethylaminomethyl-
piperidine,

1-(3,7,11,15,19-pentamethyl-2,6,10,14,18-
eicosapentaenoyl)-4-methylpiperazine,

1-(3,7,11,15,19-pentamethyl-2,6,10,14,18-
eicosapentaenoyl)-4-methyl-hexahydro-1,4-
diazepine,

N-methyl-N-(3,7,11,15,19-pentamethyl-
2,6,10,14,18-eicosapentaenoyl)-ethanolamine,

3-(3,7,11,15,19-pentamethyl-2,6,10,14,18-
eicosapentaenoylamino)-1-ethylpiperidine,

and

2-(3,7,11,15,19-pentamethyl-2,6,10,14,18-
eicosapentaenoylamino)-1-ethylpyrrolidine.

Examples of typical compounds of the second
compound group according to the invention are listed
below.

1. 2-palmitoilamino-2-ethyl-1,3-propanediol
2. 2-oleoilamino-2-ethyl-1,3-propanediol
3. 2-(3',7',11',15'-tetramethyl-hexadecanoilamino)-
1-ethyl-piperidine
4. 2-(3',7',11',15'-tetramethyl-2'-hexadecaenoilamino)-
1-ethyl-piperidine
5. 2-(3',7',11',15'-tetramethyl-hexadecanoilaminomethyl)-
1-ethyl-piroridine
6. 2-(3',7',11',15'-tetramethyl-2'-hexadecaenoilaminomethyl)-
1-ethyl-piroridine
7. N-(3,7,11,15-tetramethyl-hexadecanoilaminoethyl)-
piroridine

-
8. N-(3,7,11,15-tetramethyl-2-hexadecaenylaminoethyl)-
piperidine
9. N-(3,7,11-trimethyl-2,6,10-dodecatrienylaminoethyl)-
morpholine
- 5 10. N-(3,7,11,15-tetramethyl-hexadecanoylaminoethyl)-
morpholine
11. N-(3,7,11,15-tetramethyl-hexadecanoyl)-N',N'-
dimethyl-ethylenediamine hydrochloride
12. N-(3,7,11,15-tetramethyl-hexadecanoyl)-N',N'-
diisopropyl-ethylenediamine hydrochloride
- 10 13. N-(3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenyl)-
N',N',-diisopropyl-ethylenediamine hydrochloride
14. N-(3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenyl)-
N',N'-diethyl-ethylenediamine hydrochloride
- 15 15. N-(3,7,11,15-tetramethyl-hexadecanoyl)-N',N'-
diethyl-ethylenediamine
16. N-(3,7,11,15-trimethyl-2,6,10-dodecatrienyl)-
N',N'-diethyl-ethylenediamine
17. N'-(3,7,11-trimethyl-2,6,10-dodecatrienyl)-2-
aminopyridine
- 20 18. N'-(3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenyl)-
2-aminopyridine
19. N-(3,7,11-trimethyl-2,6,10-dodecatrienyl)-imidazol
20. N-(3,7,11,15-tetramethyl-hexadecanoyl)-imidazol
- 25 21. N-(3',7',11'-trimethyl-2',6',10',-dodecatrienyl)-
2-amino-2-enol-1,3-propanediol
22. N-(3',7',11',15'-tetramethyl-hexadecanoyl)-2-amino-
2-ethyl-1,3-propanediol
23. N-(3',7',11',15'-tetramethyl-hexadecanoyl)-3-amino-
1,2-propanediol
- 30 24. N-(3',7',11'-trimethyl-2',6',10'-dodecatrienyl)-
3-amino-1,2-propanediol

0110397

25. N-(3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenoil)-
N',N',N'',N'''-tetraethyl-diethylenetriamine
26. N-(3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenoil)-
N',N',N'',N'''-tetraisopropyl-diethylenetriamine
- 5 27. N-(3,7,11,15-tetramethyl-hexadecanoil)-N',N',N'',N'''-
tetramethyl-diethylenetriamine
28. N-(3,7,11-trimethyl-2,6,10-dodecatrienoil)-
N',N',N'',N'''-tetramethyl-diethylenetriamine
29. N-(3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenoil)-
10 N',N',N'',N'''-tetraisopropyl-diethylenetriamine hydrochloride
30. N-(3,7,11,15-tetramethyl-hexadecanoil)-N',N',N'',N'''-
tetramethyl-diethylenetriamine hydrochloride
31. N-(3,7,11-trimethyl-2,6,10-dodecatrienoil)-N',N',N'',N'''-
tetramethyl-diethylenetriamine hydrochloride
- 15 32. N-(3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenyl)-
N',N'-diethylaminomethyl-carboxamide
33. N-(3,7,11,15-tetramethyl-2-hexadecaenyl)-N',N'-
dimethylaminomethyl-carboxamide
34. N-3,7,11-trimethyl-2,6,10-dodecatrienyl)-N',N'-
20 diethylaminomethyl-carboxamide
35. N-(3,7,11,15-tetramethyl-hexadecyl)-2-aminoethyl-
carboxamide
36. N-(3,7,11-tetramethyl-2,6,10-dodecatrienyl)-2-
aminoethyl-carboxamide
- 25 37. N-(3,7,11,15-tetramethyl-hexadecyl)-N',N'-dimethyl-
ethylenediamine
38. N-(3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenyl)-
N',N'-diethyl-ethylenediamine
39. N-(4-(2',6',10'-trimethyl-1',5',9'-undecatrienyl)benzyl)
30 propanolamine
40. N-(4-(2',6'-dimethyl-1',5'-heptadienyl)-benzyl)-
ethanolamine

41. N-(4-(2',6',10'-trimethyl-1',5',9'-undecatrienyl)-benzoyl)-dipropanolamine
42. N-(4-(2',6'-dimethyl-1',5'-heptadienyl)-benzoyl)-diethanolamine
- 5 43. N-(4-(2',6',10'-trimethyl-1',8',9'-undecatrienyl)-benzoyl)-N',N'-diethylethylenediamine
44. N-(4-(2',6',10'-trimethyl-1',5',9'-undecatrienyl)-benzoyl)-N',N'-diisopropylethylenediamine
45. N-(4-(2',6',10'-trimethyl-trimethyl-1',5',9'-undecatrienyl)-benzoyl)-N',N'-diethylethylenediamine hydrochloride
- 10 46. N-(4-(2',6',10'-trimethyl-1',5',9'-undecatrienyl)-benzoyl)-N',N'-diisopropylethylenediamine hydrochloride
47. N-(4-(2',6',10'-trimethyl-1',5',9'-undecatrienyl)-benzoyl)-N',N',N'',N''-tetraethyl-diethylenetriamine
- 15 48. N-(4-(2',6'-dimethyl-1',5'-heptadienyl)-benzoyl)-N',N',N'',N''-tetramethyl-diethylenetriamine
49. N-(4-(2',6',10'-trimethyl-1',5',9'-undecatrienyl)-benzoyl)-N',N',N'',N''-tetraethyl-diethylenetriamine hydrochloride
- 20 50. N-(4-(2',6',10'-trimethyl-undecyl)-benzoyl)-propanolamine
51. N-(4-(2',6'-dimethyl-heptyl)-benzoyl)-diethanolamine
52. N-(4-(2',6',10'-trimethyl-undecyl)-benzoyl)-N',N'-diethylethylenediamine
53. N-(4-(2',6',10'-trimethyl-undecyl)-benzoyl)-N',N'-diethylethylenediamine hydrochloride
- 25 54. N-(4-(2',6',10'-trimethyl-undecyl)-benzoyl)-N',N',N'',N''-tetraethyl-diethylenetriamine
55. N-(4-(2',6',10'-trimethyl-undecyl)-benzoyl)-N',N',N'',N''-tetraethyl-diethylenetriamine hydrochloride
- 30 56. N-(4-(2',6'-dimethyl-1',5'-heptadienyl)-benzoyl)ethanolamine
57. N-(4-(2',6'-dimethylheptyl)-benzoyl)propanolamine
58. N-(4-(2',6'-dimethyl-1',5'-heptadienyl)-benzoyl)-N',N',N'',N''-tetramethyl-diethylenetriamine

59. N-(4-(2',6'-dimethyl-1',5'-heptadienyl)-benzoyl)-
N',N',N'',N''-tetraethyl-diethylenetriamine
60. N-(4-(2',6'-dimethylheptyl)-benzoyl)-N',N',N'',N''-
tetramethyl-diethylenetriamine hydrochloride
- 5 61. N-(4-(2',6',10',14'-tetramethyl-1',5',9',13'-pentadeca-
tetraenyl)-benzoyl)-ethanolamine
62. N-(4-(2',6',10',14'-tetramethyl-pentadecyl)-benzoyl)-
propanolamine
63. N-(4-(2',6',10',14'-tetramethyl-1',5',9',13'-pentadecatetra-
10 enyl)-benzoyl)-N',N',N'',N''-tetramethyl-diethylenetriamine
64. N-(3-(4'-(4'',8''-dimethyl-3'',7''-nonadienyl)phenyl)-
butanoyl)ethanolamine
65. N-(3-(4'-(4'',8''-dimethyl-3'',7''-nonadienyl)phenyl)-
butanoyl)-N',N',N'',N''-tetramethyl-diethylenetriamine
- 15 66. N-(3-(4'-(4'',8''-dimethyl-3'',7''-nonadienyl)-phenyl)-
2-butenoyl)ethanolamine
67. N-(3-(4'-(4'',8''-dimethyl-3'',7''-nonadienyl)-phenyl)-
2-butenoyl)-N',N',N'',N''-tetramethyl-diethylenetriamine
68. N-(2-methyl-4-(2',6',10'-trimethyl-1',5',9'-undecatrienyl)-
20 benzoyl)ethanolamine
69. N-(2-methyl-4-(2',6',10'-trimethyl-1',5',9'-undecatrienyl)-
benzoyl)-N',N',N'',N''-tetramethyl-diethylenetriamine
70. N-(2-floro-4-(2',6',10'-trimethyl-1',5',9'-undecatrienyl)-
benzoyl)-ethanolamine

25

30

0110397

71. N-(2-floro-4-(2',6',10'-trimethyl-1',5',9'-undecatrienyl)-benzoyl)-N',N',N'',N''-tetramethyl-diethylenetriamine
72. N-(4-(2',6',10'-trimethyl-1',5',9'-undecatrienyl)-1-naphthoyl)-ethanolamine
- 5 73. N-(4-(2',6',10'-trimethyl-1',5',9'-undecatrienyl)-1-naphthoyl)-N',N',N'',N''-tetraethyl-diethylenetriamine
74. N-(5-(2',6',10'-trimethyl-1',5',9'-undecatrienyl)-1-naphthoyl)ethanolamine
75. N-(5-(2',6',10'-trimethyl-1',5',9'-undecatrienyl)-1-naphthoyl)-N',N',N'',N''-tetramethyl-diethylenetriamine
- 10 76. N-(4-(2',6',10'-trimethyl-undecyl)-5,6,7,8-tetrahydro-1-naphthoyl)propanolamine
77. N-(4-(2',6',10'-trimethyl-undecyl)-5,6,7,8-tetrahydro-1-naphthoyl)-N',N',N'',N''-tetraethyl-diethylenetriamine
- 15 78. N-(4-(2',6',10'-trimethyl-1',5',9'-undecatrienyl)-5,6,7,8-tetrahydro-1-naphthoyl)ethanolamine
79. N-(4-(2',6',10'-trimethyl-1',5',9'-undecatrienyl)-5,6,7,8-tetrahydro-1-naphthoyl)-N',N',N'',N''-tetramethyl-diethylenetriamine

20

25

30

The polyprenyl compound according to the invention have excellent anti-PAF and antithrombic activities, and are useful in pharmaceutical compositions intended for utilizing these anti-PAF and antithrombic activities. The term "PAF" means platelet activating factor. Barbaro et al found in 1966 that a rabbit basophile sensitized by immunoglobulin E (IgE) released a factor which caused platelet degranulation and aggregation. This factor was named PAF by Benveniste et al in 1972. Demopoulos et al reported in 1979 that its structure was identical with that of 1-alkyl-2-acetyl-sn-glycero-3-phosphocholine. PAF is an alkyl ether phospholipid having an acetyl group which is a new mediator for platelet aggregation.


Human disseminated intravascular coagulation syndrome (DIC) is a condition wherein blood coagulation is abnormally promoted, blood in the microcardiovascular system is widely coagulated and many thrombi are formed. One of the factors causing this condition is thrombin. As noted above, the compounds of the present invention have antithrombic and anti-PAF activity. Therefore, the compounds of the present invention are useful as excellent antithrombic drugs having both antithrombic and anti-PAF activities for the treatment of DIC.

The formation of thrombi can further cause hemadostenosis, angiostenosis and ischemic lesions or infarctions in principal internal organs such as heart, brain and lungs. Therefore, the compounds of the present invention are useful for the therapy and prophylaxis of myocardial angina pectoris, cerebral thrombosis, DIC and chronic arteriosclerosis. In addition to being useful as drugs for the therapy and prophylaxis of these thromboses, the compounds of the present invention having anti-PAF activity can be used as anti-inflammatory drugs, antiasthmatic drugs, antiarteriosclerotic drugs, antishock and blood pressure controlling drugs, immune function controlling drugs and antiallergic drugs.

0110397

We have unexpectedly found that the compounds of the present invention have anti-PAF and antithrombic activities. The present invention is based on this finding. No compounds similar in structure to those of the present invention have been known to exhibit an anti-PAF activity.

The first compound group according to the invention wherein W in the formula (I) is the group of -COR can be produced by various methods. For example a polyprenyl carboxylic acid of the formula (II)



wherein A, B, Y, Z and n have the same meanings as defined above, or a reactive derivative thereof, is reacted with an amine of the formula RH [III], wherein R has the same meaning as defined above, to amidate said carboxylic acid or its derivative, thus producing the desired polyprenylcarboxylic acid amide [I]. Examples of reactive derivatives of the carboxylic acid [II] include halides, anhydrides and mixed anhydrides of the acid [II].

If desired, the reaction can be carried out in the presence of a dehydrating agent such as N,N'-dicyclohexylcarbodiimide, N,N'-diethylcarbodiimide, trialkyl phosphate, ethyl polyphosphate or tosyl chloride, in order to conduct the reaction smoothly.


Further, in order to capture the hydrogen halide formed by the reaction and promote the reaction, a base can be added. Examples of such bases include inorganic bases, such as potassium hydroxide, sodium hydroxide, potassium carbonate and sodium carbonate, and tertiary amines, such as pyridine and triethylamine. The reaction can ordinarily be conducted in a solvent, such as dioxane, tetrahydrofuran, dimethyl sulfoxide or a lower alcohol, or mixtures thereof.

0110397





The second compound group according to the invention wherein W in the formula (I) is X can be produced by various preparation methods. Some examples therefor are described below.

- 5 (1) When X is one of the groups (6) to (14) defined hereinbefore, the following preparation may be used.





The intended product can be obtained by reacting a polyprenyl carboxylic acid of the
10 formula (II)




15 wherein A, B, Y and Z are defined hereinbefore and n is an integer of 1 to 3, or a reactive derivative thereof, with one of amine compounds of the formulae:



0110397




and




Also in these preparation methods, the reactive derivative, a dehydrating agent, a base and a solvent may be used in the same manner as described hereinbefore.

(2) When X is the group (15), the product can be obtained by reacting a polyprenyl compound of the formula (IV)




wherein A, B, Y, Z and n are defined hereinbefore, with a carboxylic acid of the formula (V)

0110397




in the same manner as shown in the before metioned
(1) to carry our amidation.


(3) When X is the group (16), the product can
be obtained by reacting a halide of a polyprenyl
compound of the formula (VI)




wherein Hal is a halogen and A, B, Y, Z and n are
defined hereinbefore, with an amine compound of
the formula (VII) in order to effect dehalogenation.



(4) When X is the group (1), the product can
be obtained by reacting a polyprenyl compound of
the formula (VIII).




with, for example, a thionyl chloride in order to
produce an acid halide of the formula (IX)




and then reacting the resulting acid halide with an amine compound of the formula (X): $R^{22}H$ in order to produce the intended product.

- (5) When X is the group (2), the intended product
 5 can be obtained by reacting a polyprenyl compound of the formula (XI)






with, for example, thionyl chloride to give an acid halide of the formula (XII)



wherein Hal is a halogen,

- 20 and then reacting the acid halide with an amine compound of the formula (X): $R^{22}H$.

- (6) When X is one of the groups (3), (4) and
 (5), the product can be obtained by reacting a polyprenyl
 compound of each of the formulae (XIV), (XV) and
 25 (XVI)



0110397

with, for example, thionyl chloride in order to give an acid halide of each polyprenyl compound and then reacting the resultant with an amine compound of the formula (X).

5 In the preparation methods as mentioned above, oxalyl chloride may be placed for thionyl chloride.

10 The following experimental example is provided to illustrate the effects of the compounds according to the invention.

15

20

25

30

Experimental Example

(1) Anti-PAF activity

Experimental method

(a) Preparation of washed platelet suspension
(hereinafter referred to as W.P.)

5 A blood sample was collected from a carotid artery of a male rabbit weighing 2.5 kg, while adding thereto one volume of a 3.13% sodium citrate solution per 9 volumes of blood as a coagulant. The resulting blood
10 was centrifuged at 200 xg for 20 minutes to separate out platelet-rich plasma (hereinafter referred to as PRP). This PRP was centrifuged at 1,000 xg for 15 minutes to separate the platelets from the plasma. The deposited platelets were washed twice with a Tyrode solution
15 (Tyrode- Ca^{++}), from which Ca^{++} had been removed and to which 1% bovine serum albumin (BSA) had been added, such that 9×10^5 platelets per μl were finally suspended in the Tyrode solution.

(b) Preparation of specimens and PAF






20 Specimens were dissolved or suspended in a physiological saline. PAF was dissolved in a Tyrode solution containing 1% BSA. PAF synthesized from D-mannitol in accordance with a method disclosed by J.J. Godfroid et al was used (FEBS Letters 116, 161-164,
25 1980).

(c) Measurement of platelet aggregation

Platelet aggregation measurements were conducted according to nephelometry described by Born et al using a platelet aggregation meter manufactured by Schencko Co.
30 Specimen solutions having various concentrations of test compounds and 0.25 ml of W.P. were subjected to preincubation at 37°C for 4 minutes, and then PAF was added thereto to give a final PAF concentration of 30 ng/ml to provoke platelet aggregation. The

0110397

Table 1 - Continued

Compound	Concent- ration (μ M)	Inhibition rate PAF (%)	Inhibition rate Thrombin (%)
	50	51.2	43.3
	50	93.1	83.7
	50	100	92
	50	93.9	86.5
	50	76.4	—

0110397

Table 1 - Continued

Compound	Concent- ration (μ M)	Inhibition rate P _{AF} (%)	Inhibition rate Thrombin (%)
$\text{H} \left(\text{CH}_2 \right)_2 \text{CH} = \text{CH} \left(\text{CH}_2 \right)_2 \text{C}(=\text{O})\text{NHCH}_2\text{CH}_2\text{OH}$	50	31.4	—
$\text{H} \left(\text{CH}_2 \right)_4 \text{CH} = \text{CH} \left(\text{CH}_2 \right)_2 \text{C}(=\text{O})\text{NHCH}_2\text{CH}_2\text{OH}$	50	22.7	—
$\text{H} \left(\text{CH}_2 \right)_2 \text{CH} = \text{CH} \left(\text{CH}_2 \right)_2 \text{C}(=\text{O})\text{NHCH}_2\text{CH}_2\text{OH}$	50	18.7	—
$\text{H} \left(\text{CH}_2 \right)_3 \text{CH} = \text{CH} \left(\text{CH}_2 \right)_2 \text{C}(=\text{O})\text{N} \begin{array}{c} \diagup \text{OH} \\ \diagdown \end{array}$	50	27.4	19.2
$\text{H} \left(\text{CH}_2 \right)_3 \text{CH} = \text{CH} \left(\text{CH}_2 \right)_2 \text{C}(=\text{O})\text{N} \begin{array}{c} \diagup \text{CH}_2\text{OH} \\ \diagdown \end{array}$	50	34.4	27.4

0110397

transmittance of the W.P. before aggregation, that is,
before the addition of the aggregation provoking agent,
was rated as 0, and the transmittance of the Tyrode
solution was rated as 100. After the addition of the
5 PAF solution to the W.P., light transmittance increased
as aggregation proceeded. The value of the light
transmittance at the time when the aggregation had
proceeded to a maximum was rated as the maximum
aggregation (hereinafter referred to as MA) which was
10 used as an index of the degree of aggregation.

The inhibition rate was calculated according to the
following equation. The aggregation in a W.P.-containing
physiological saline solution free of test compound, as
a control, was rated as 0% aggregation inhibition.

$$15 \quad \text{Aggregation inhibition rate (\%)} \\ = \frac{\text{M.A. Control} - \text{M.A. Sample}}{\text{M.A. Control}} \times 100$$

wherein

20 M.A. Control: maximum aggregation after platelet
aggregation in PAF was provoked,
after the addition of physiological
saline.

M.A. Sample: maximum aggregation after platelet
aggregation in PAF was provoked,
after the addition of the test
25 compound.

(2) Antithrombic activity

The measurement of the antithrombic activity was
conducted in the same manner as described above, except
that bovine thrombin at a final concentration of 0.2
30 units/ml was used as the platelet aggregation provoking
agent in place of PAF.

The results are shown in Table 1.

0110397

Table 1

Compound	Concentration (μM)	Inhibition rate PAF (%)	Inhibition rate Thrombin (%)
$\text{H} \left(\text{CH}_2 \right)_3 \text{CH} = \text{CH} \text{CH}_2 \text{CH}_2 \text{C}(=\text{O})\text{NHCH}_2\text{CH}_2\text{OH}$	50	50	—
$\text{H} \left(\text{CH}_2 \right)_3 \text{CH} = \text{CH} \text{CH}_2 \text{CH}_2 \text{C}(=\text{O})\text{NHCH}_2\text{CH}_2\text{CH}_2\text{OH}$	50	22.1	11.6
$\text{H} \left(\text{CH}_2 \right)_3 \text{CH} = \text{CH} \text{CH}_2 \text{CH}_2 \text{C}(=\text{O})\text{N}(\text{OH})\text{CH}_2\text{CH}_2\text{OH}$	50	16.3	3.9
$\text{H} \left(\text{CH}_2 \right)_3 \text{CH} = \text{CH} \text{CH}_2 \text{CH}_2 \text{C}(=\text{O})\text{NHCH}_2\text{CH}_2\text{NH}_2$	50	80.8	—
$\text{H} \left(\text{CH}_2 \right)_3 \text{CH} = \text{CH} \text{CH}_2 \text{CH}_2 \text{C}(=\text{O})\text{NHCH}_2\text{CH}_2\text{N}(\text{OH})_2$	50	90.8	70.8






0110397

Table 1 - Continued

Compound	Concentration (μM)	Inhibition rate μAr (%)	Inhibition rate Thrombin (%)
	50	80.3	24.2
	50	87.0	79.6



0110397

Table 1 - Continued

Compound	Concentration (μ M)	Inhibition rate PAF (%)	Inhibition rate Thrombin (%)
	20	57.8	
	20	59.4	
	20 50	57.8 82.6	
	20	55.9	
	20 50	63.2 82.2	






0110397

Table 1 - Continued

Compound	Concentration (μ M)	Inhibition rate PAF (%)	Inhibition rate Thrombin (%)
	20	50.0	
	20	61.4	

0110397

Table 1 - Continued

Compound	Concentration (μ M)	Inhibition rate PAF (%)	Inhibition rate Thrombin (%)
 <chem>CCCCCCCCCCCCCCCC(=O)c1ccc(F)cc1N(C)C</chem>	20	64.9	
 <chem>CCCCCCCCCCCCCCCC(=O)c1cccc2ccccc12N(C)C</chem>	20	59.3	
 <chem>CCCCCCCCCCCCCCCC(=O)c1cccc2ccccc12N(C)C</chem>	20	58.3	
 <chem>CCCCCCCCCCCCCCCC(=O)c1cccc2ccccc12N(C)C</chem>	20	50.5	
 <chem>CCCCCCCCCCCCCCCC(=O)c1cccc2ccccc12N(C)C</chem>	20	60.4	




0110397

Table 1 - Continued

Compound	Concentration (μM)	Inhibition rate PAF (%)	Inhibition rate Thrombin (%)
	20	59.1	
	20	37.1	
	20	74.3	
	20	88.7	76.4
	20	73.5	34.2

0110397

Table 1 - Continued

Compound	Concentration (μ M)	Inhibition rate PAF (%)	Inhibition rate Thrombin (%)
 $\cdot \text{HCl}$	20	80.6	70.3
 $\cdot 2\text{HCl}$	20	93.6	70.6
 $\cdot 2\text{HCl}$	20	74.7	78.8

It is apparent from the above experimental results that the compounds of the present invention have an excellent anti-PAF activity. In addition to being useful as drugs for the prophylaxis and therapy of thromboses, the compounds of the invention are also
5 useful as anti-inflammatory drugs, antiasthmatic drugs, antiarteriosclerotic drugs, antishock and blood pressure controlling drugs, immune function controlling drugs and antiallergic drugs.

10 The compounds of the present invention have a very low toxicity and high safety, and are suitable for long-term continuous administration. The present invention is very valuable in this sense. When the compounds of the present invention described in the
15 above experimental example were orally administered in doses of 500 mg/kg to SD rats (each weighing about 200 g), no deaths or side effects were observed.

Dosages of the compounds of the present invention administered as a drug exhibiting an anti-PAF activity
20 to human or animal patients vary greatly depending on the type and extent of the disease, the particular compound employed and the age of the patients, and, thus, the dosage amount is not particularly limited. Generally, however, the compounds of the invention are
25 orally or parenterally administered at dosages in the range of from 10 to 1,000 mg/day/adult, preferably about 50 to 300 mg/day/adult. The unit dosage forms of drugs to be administered include powders, fine-grained
30 powders, granules, tablets, capsules and injection liquids. Such drug forms are prepared by conventional methods using conventional pharmaceutical carriers.

In the formulation of solid preparations for oral administration, an excipient is added to a base. If

desired, a binder, disintegrator, lubricant, colorant, flavoring agent, and other conventional additives are added thereto. The mixture is then shaped into powder, coated powder, granules or capsules by any conventional method.

Examples of suitable excipients are lactose, corn starch, sucrose, glucose, sorbitol, crystalline cellulose and silicon dioxide. Examples of suitable binders are polyvinyl alcohol, polyvinyl ether, ethylcellulose, methylcellulose, acacia, tragacanth, gelatin, shellac, hydroxypropylcellulose, hydroxypropylstarch and polyvinylpyrrolidone. Examples of disintegrators are starch, agar-agar, gelatin powder, crystalline cellulose, calcium carbonate, sodium hydrogencarbonate, calcium citrate, dextrin and pectin. Examples of lubricants include magnesium stearate, talc, polyethylene glycol, silica and hardened vegetable oil. As colorants, any pharmaceutically acceptable substances may be used. Examples of flavoring agents are cocoa powder, menthol, peppermint oil, borneol and cinnamon powder. The tablets and granules can be coated with sugar, gelatin or other coating agents.

In the formulation of an injection liquid, a pH adjuster, a buffer, a stabilizer, a solubilizer, etc. are added to a base to form a preparation for hypodermic, intramuscular or intravenous injection by any conventional method.

The compounds of the present invention can be administered orally or parenterally to animals such as domestic animals and poultry. Oral administration can usually be conducted by blending the compounds with ordinary feed. In parenteral administration, an injection is prepared by a conventional method and the

0110397

compound is hypodermically administered, intramuscularly or intravenously, to the animal.

The following is an example of a preparation containing N-(3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenoyl)-ethanolamine, hereinafter referred to as the base, a typical compound of the present invention, as the active ingredient.

Formulation of preparation (tablet)

	base	10 g
10	silicic anhydride	50 g
	crystalline cellulose	70 g
	corn starch	36 g
	hydroxypropylcellulose	10 g
	magnesium stearate	4 g

15 The above ingredients are formulated into tablets (180 mg per tablet) by a conventional method.

The following examples of preparation of compounds according to the invention are provided to illustrate the present invention, but are not to be construed as
20 limiting the invention in any way.

Example 1

N-(3,7,11,15-Tetramethyl-2,6,10,14-hexadecatetraenoyl)-ethanolamine

25 6.1 g of 3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenoic acid was dissolved in 50 ml of tetrahydrofuran and 3.1 ml of triethylamine was added thereto. While cooling the mixture in ice with stirring, 2.1 ml of ethyl chlorocarbonate was added dropwise. Then the mixture was stirred for 15 minutes
30 and 1.8 ml of ethanolamine was added thereto. After the mixture was stirred for 30 minutes at room temperature, water was added thereto and the resulting aqueous

0110397

solution was extracted with ethyl acetate. The ethyl acetate layer was separated from the aqueous layer, washed with 5% aqueous hydrochloric acid solution and then water, and dried over magnesium sulfate. The solvent was distilled off. The resulting reaction mixture was subjected to chromatography on a silica gel column to afford 6.5 g (yield 94%) of the title compound as a colorless oil.

Elemental analysis for $C_{22}H_{37}NO_2$

10		C	H	N
	calculated (%)	76.03	10.73	4.03
	found (%)	76.00	10.31	3.94

° Mass (m/z) : 347 (M^+)

15 ° NMR (δ , $CDCl_3$) : 1.59 (9H,s), 1.68 (3H,s)
 1.9 - 2.2 (12H), 2.12 (3H,d,J=1),
 2.90 (1H,br), 3.3 - 3.5 (2H), 3.35 - 3.7 (2H),
 5.06 (3H,m), 5.52 (1H,s), 5.94 (1H,br,s).

Example 2

N-(3,7,11,15-Tetramethyl-2-hexadecenoyl)-ethanolamine

20 The procedure of Example 1 was repeated except that 6.2 g of 3,7,11,15-tetramethyl-2-hexadecenoic acid and 1.8 ml of ethanolamine were used as starting materials. 6.7 g (yield 95%) of the title compound was obtained as
 25 a colorless oil.

0110397

Elemental analysis for $C_{22}H_{43}NO_2$

	C	H	N
calculated (%)	74.73	12.26	3.96
5 found (%)	74.54	12.33	3.88

° Mass (m/z) : 353 (M^+)° NMR (δ , $CDCl_3$) : 0.84 (12H,d,J=7),

1.0 - 1.5 (20H,m), 1.81 (3H,d,J=1),

10 2.4 - 2.7 (2H,m), 3.3 - 3.5 (2H,m),

3.6 - 3.8 (2H,m), 5.52 (1H,s), 5.70 (1H,br).

Example 3

15 N-(3,7,11,15-Tetramethyl-hexadecanoyl)-
ethanolamine

6.5 g of N-(3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenoyl)-ethanolamine obtained in Example 1 was hydrogenated in the presence of a catalyst composed of palladium on carbon in 40 ml of ethanol. The ethanol
 20 layer was then separated from the catalyst and the solvent was distilled off to afford 6.7 g (yield 94%) of the title compound as a colorless oil.

Elemental analysis for $C_{22}H_{45}NO_2$

	C	H	N
25 calculated (%)	74.30	12.76	3.94
found (%)	74.20	12.84	3.91

° Mass (m/z) : 355 (M^+)30 ° NMR (δ , $CDCl_3$) : 0.86 (15H,d,J=6),

1.0 - 1.5 (23H), 1.96 (2H,m),

3.3 - 3.8 (4H), 6.05 (1H,br).

Example 4N-(3,7,11,15-Tetramethyl-6,10,14-hexadecatrienoyl)-ethanolamine

5 The procedure of Example 1 was repeated except that 6.1 g of 3,7,11,15-tetramethyl-6,10,14-hexadecatrienoic acid and 1.8 ml of ethanolamine were used as starting materials. 6.4 g (yield 92%) of the title compound was obtained as a colorless oil.

10 Elemental analysis for $C_{22}H_{39}NO_2$

	C	H	N
calculated (%)	75.59	11.25	4.01
found (%)	75.47	11.39	4.08

15 ° Mass (m/z) : 349 (M^+)

° NMR (δ , $CDCl_3$) : 0.94 (3H,d,J=5),

1.1 - 1.5 (3H,m), 1.60 (9H,s), 1.68 (3H,s),

1.8 - 2.2 (12H), 3.3 - 3.8 (4H,m),

4.16 (1H,br), 5.09 (3H,m), 6.72 (1H,br).

20

Example 5N-(3,7,11,15,19-Pentamethyl-2,6,10,14,18-eicosapentaenoyl)-ethanolamine

25 The procedure of Example 1 was repeated except that 7.4 g of 3,7,11,15,19-pentamethyl-2,6,10,14,18-eicosapentaenoic acid and 1.8 ml of ethanolamine were used as starting materials. 7.7 g (yield 93%) of the title compound as a colorless oil was obtained.

30 Elemental analysis for $C_{27}H_{45}NO_2$

	C	H	N
calculated (%)	78.02	10.91	3.37
found (%)	77.93	10.99	3.30

0110397

- ° Mass (m/z) : 415 (M^+)
 ° NMR (δ , $CDCl_3$) : 1.60 (12H,s), 1.68 (3H,s),
 1.8 - 2.2 (17H), 2.14 (3H,d,J=1),
 3.2 - 3.8 (4H), 5.08 (4H,m), 5.59 (1H,br,s),
 6.26 (1H,br,t).

Example 6N-(3,7,11-Trimethyl-2,6,10-dodecatrienoyl)-ethanolamine

The procedure of Example 1 was repeated except that 4.7 g of 3,7,11-trimethyl-2,6,10-dodecatrienic acid and 1.8 ml of ethanolamine were used as starting materials. 5.2 g (yield 94%) of the title compound was obtained as a colorless oil.

Elemental analysis for $C_{17}H_{29}NO_2$

	C	H	N
calculated (%)	73.07	10.46	5.01
found (%)	73.00	10.53	5.06

- ° Mass (m/z) : 279 (M^+)
 ° NMR (δ , $CDCl_3$) : 1.60 (6H,s), 1.68 (3H,s),
 1.8 - 2.3 (8H), 2.12 (3H,s), 3.2 - 3.8 (4H),
 4.20 (1H,br), 5.08 (2H,m), 5.60 (1H,br,s),
 6.76 (1H,br).

Example 7N-(3,7,11,15-Tetramethyl-2,6,10,14-hexadecatetraenoyl)-diethanolamine

The procedure of Example 1 was repeated except that 6.1 g of 3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenoic acid and 2.9 ml of diethanolamine

0110397

were used as starting materials. 7.0 g (yield 90%) of the title compound was obtained as a colorless oil.

Elemental analysis for $C_{24}H_{41}NO_3$

	C	H	N
5 calculated (%)	73.61	10.55	3.58
found (%)	73.52	10.66	3.51

° Mass (m/z) : 391 (M^+)

10 ° NMR (δ , $CDCl_3$) : 1.61 (9H,s), 1.68 (3H,s),
 1.8 - 2.2 (12H), 2.12 (3H,d,J=2),
 3.4 - 3.9 (8H), 4.56 (2H,br), 5.08 (3H,m),
 5.88 (1H,br,s).

15 Example 8

N-(3,7,11,15-Tetramethyl-2,6,10,14-hexadecatetraenoyl)-glycine

The reaction of 6.1 g of 3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenoic acid with glycine
 20 ethyl ester hydrochloride was conducted in the same
 manner as described in Example 1 except that 6.5 ml of
 triethylamine was used. Then, a solution of 2.7 g of
 potassium hydroxide in ethanol was added. The mixture
 was heated under reflux for 30 minutes. After the
 25 completion of the reaction, water was added and the
 obtained aqueous solution was extracted with ethyl
 acetate. The ethyl acetate layer was washed with water
 and dried over magnesium sulfate. The solvent was
 distilled off to afford 6.1 g (yield 85%) of the title
 30 compound as a pale brown oil.

0110397

Elemental analysis for $C_{22}H_{35}NO_3$

	C	H	N
calculated (%)	73.09	9.76	3.87
5 found (%)	72.97	9.80	3.79

° Mass (m/z) : 361 (M^+)

° NMR (δ , $CDCl_3$) : 1.59 (9H,s), 1.66 (3H,s),
 1.7 - 2.2 (12H), 2.10 (3H,s), 4.02 (2H,br,
 10 d,J=5), 5.04 (3H,m), 5.60 (1H,br,s),
 6.08 (1H,br), 6.37 (1H,br).

Example 9

15 N-(3,7,11,15-Tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-ethylenediamine

3.1 ml of triethylamine was added to a solution of
 6.1 g of 3,7,11,15-tetramethyl-2,6,10,14-
 hexadecatetraenoic acid in 50 ml of tetrahydrofuran.
 While cooling the mixture in ice with stirring, 2.1 ml
 20 of ethyl chlorocarbonate was added dropwise and the
 mixture was stirred for 15 minutes. Then 2.0 ml of
 ethylenediamine was added thereto and the mixture was
 stirred for 30 minutes at room temperature. Then water
 was added thereto and the resulting solution was
 25 extracted with chloroform.

The chloroform layer was washed with water and
 dried over magnesium sulfate. The solvent was distilled
 off. The resulting reaction mixture was chromatographed
 on a silica gel column to afford 5.5 g (yield 80%) of
 30 the title compound as a pale brown oil.

0110397

Elemental analysis for $C_{22}H_{38}N_2O$

	C	H	N
calculated (%)	76.25	11.05	8.08
5 found (%)	76.22	11.10	8.10

° Mass (m/z) : 346 (M^+)° NMR (δ , $CDCl_3$) : 1.56 (9H,s), 5.05 (3H,m),

1.64 (3H,s), 5.56 (1H,br,s), 1.75 - 2.2 (14H),

10 6.48 (1H,br), 2.16 (3H,d,J=1),

2.7 - 2.95 (2H), 3.1 - 3.4 (2H).

Example 10

15 N-(3,7,11,15-Tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-ethylenediamine hydrochloride

Hydrogen chloride gas was passed through a methanol
 solution of N-(3,7,11,15-tetramethyl-2,6,10,14-
 hexadecatetraenoyl)-ethylenediamine obtained in Example
 9. The solvent was distilled off to afford 6.1 g of the
 20 title compound as a brown oil.

Elemental analysis for $C_{22}H_{39}N_2OCl$

	C	H	N	Cl
calculated (%)	68.99	10.26	7.32	9.26
25 found (%)	68.79	10.50	7.15	9.20

° Mass (m/z) : 384 (M^+ , Cl^{37}) 382 (M^+ , Cl^{35})° NMR (δ , $CDCl_3$) : 1.60 (9H,s), 1.68 (3H,s),

1.7 - 2.3 (15H), 2.18 (3H,d,J=1),

30 3.0 - 4.0 (4H), 5.08 (3H,m), 5.74 (1H,br),

7.80 (1H,br).

Example 11N-(3,7,11,15-Tetramethyl-2,6,10,14-hexadecatetraenoyl)-N',N'-dimethylethylenediamine

The procedure of Example 9 was repeated except that
 5 6.1 g of 3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenoic acid and 3.3 ml of N,N-dimethylethylene diamine were used as starting materials. 6.5 g (yield 88%) of the title compound was obtained as a pale yellow oil.

10 Elemental analysis for $C_{24}H_{42}N_2O$

	C	H	N
calculated (%)	76.95	11.30	7.48
found (%)	76.85	11.31	7.43

15

° Mass (m/z) : 374 (M^+)

° NMR (δ , $CDCl_3$) : 1.60 (9H,s), 1.68 (3H,s),

1.8 - 2.2 (12H), 2.24 (6H,s), 2.26 (3H,s),

2.3 - 2.5 (2H), 2.8 - 3.0 (2H), 5.10 (3H,m),

20

5.56 (1H,br,s), 6.14 (1H,br).

Example 12N-(3,7,11,15-Tetramethyl-2,6,10,14-hexadecatetraenoyl)-N',N',N'-trimethylethylenediamine chloride

25

Chloromethane gas was passed through a solution of
 6.5 g of N-(3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenoyl)-N',N'-dimethylethylenediamine
 obtained in Example 11, dissolved in 50 ml of benzene.

30

The solvent was distilled off to afford 7.4 g of the title compound as a white wax.

0110397

Elemental analysis for $C_{25}H_{45}N_2OCl$

	C	H	N	Cl
calculated (%)	70.63	10.67	6.57	8.34
5 found (%)	70.69	10.51	6.58	8.19

° Mass (m/z) : 426 (M^+ , Cl^{37}), 424 (M^+ , Cl^{35})

° NMR (δ , $CDCl_3$) : 1.60 (9H,s), 1.68 (3H,s),
 1.8 - 2.2 (12H), 2.14 (3H,d,J=2), 2.24 (9H,s),
 10 2.3 - 2.5 (2H), 3.2 - 3.5 (2H), 5.08 (3H,m),
 5.56 (1H,br,s), 6.04 (1H,br).

Example 13

15 N-(3,7,11,15-Tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-N',N',N'-trimethylethylenediamine iodide

3.4 g of methyl iodide was added to 6.5 g of
 N-(3,7,11,15-tetramethyl-2,6,10,14-hexadecatetra-
 enoyl)-N',N'-dimethylethylenediamine obtained in Example
 11. The mixture was left to stand at room temperature
 20 for 15 minutes. An excess amount of methyl iodide was
 distilled off to afford 9.0 g of the title compound as a
 brown solid with a melting point in the range of 53 to
 55°C.

Elemental analysis for $C_{25}H_{45}N_2OI$

25

	C	H	N	I
calculated (%)	58.17	8.78	5.42	24.57
found (%)	57.98	8.80	5.41	24.66

30

° Mass (m/z) : 516 (M^+)

° NMR (δ , $CDCl_3$) : 1.60 (9H,s), 1.68 (3H,s),
 1.8 - 2.2 (12H), 2.16 (3H,d,J=2), 3.46 (9H,s),
 3.84 (4H,br,s), 5.08 (3H,m), 5.72 (1H,br,s),
 7.40 (1H,br).

Example 14N-Methyl-N-(3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenoyl)-ethanolamine

The procedure of Example 1 was repeated except that 6.1 g of 3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenoic acid and 2.4 ml of N-methyl-ethanolamine were used. 6.6 g (yield 92%) of the title compound was obtained as a colorless oil.

Elemental analysis for $C_{23}H_{39}NO_2$

	C	H	N
calculated (%)	76.40	10.87	3.87
found (%)	76.38	10.90	3.90

° Mass (m/z) : 361 (M^+)

° NMR (δ , $CDCl_3$) : 1.58 (9H,s), 1.64 (3H,s),
1.8 - 2.2 (12H), 2.12 (3H,s), 2.95 (3H,d,J=1),
3.2 - 3.8 (4H), 4.30 (1H,br), 5.06 (3H,m),
5.76 (1H,br,s).

Example 15N-(3,7,11,15-Tetramethyl-2,6,10,14-hexadecatetraenoyl)-3-hydroxypiperidine

The procedure of Example 1 was repeated except that 6.1 g of 3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenoic acid and 3.0 g of 3-hydroxypiperidine were used as starting materials. 7.4 g (yield 96%) of the title compound was obtained as a colorless oil.

Elemental analysis for $C_{25}H_{41}NO_2$

0110397

	C	H	N
calculated (%)	77.47	10.67	3.61
found (%)	77.41	10.71	3.59

- 5 ° Mass (m/z) : 387 (M^+)
 ° NMR (δ , $CDCl_3$) : 1.60 (9H,s), 1.68 (3H,s),
 1.6 - 2.2 (16H), 2.12 (3H,br,s),
 2.7 - 3.9 (6H), 5.08 (3H,m),
 5.82 (1H,br,s).

10

Example 16N-(3,7,11,15-Tetramethyl-2,5,10,14-hexadeca-
tetraenoyl)-2-hydroxymethylpiperidine

- The procedure of Example 1 was repeated except that
 15 6.1 g of 3,7,11,15-tetramethyl-2,6,10,14-
 hexadecatetraenoic acid and 3.5 g of 2-hydroxy-
 methylpiperidine were used as starting materials. 7.5 g
 (yield 94%) of the title compound was obtained as a
 colorless oil.

- 20 Elemental analysis for $C_{26}H_{43}NO_2$

	C	H	N
calculated (%)	77.75	10.79	3.49
found (%)	77.73	10.83	3.30

25

- ° Mass (m/z) : 401 (M^+)
 ° NMR (δ , $CDCl_3$) : 1.60 (9H,s), 1.68 (3H,s),
 1.7 - 2.2 (18H), 2.12 (3H,br,s),
 3.2 - 4.0 (6H), 5.08 (3H,m), 5.74 (1H,br).

30

Example 17N-(3,7,11,15-Tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-propanolamine

0110397

The procedure of Example 1 was repeated except that 6.1 g of 3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenoic acid and 2.3 ml of propanolamine were used as starting materials. 6.6 g (yield 92%) of the title compound was obtained as a colorless oil.

Elemental analysis for $C_{23}H_{39}NO_2$

	C	H	N
calculated (%)	76.40	10.87	3.87
10 found (%)	76.23	10.95	3.77

° Mass (m/z) : 361 (M^+)

° NMR (δ , $CDCl_3$) : 1.59 (9H,s), 1.67 (3H,s),

1.5 - 1.8 (2H), 2.13 (3H,d,J=1),

15 1.9 - 2.2 (13H), 3.3 - 3.7 (4H), 5.09 (3H,m),

5.56 (1H,br,s), 6.03 (1H,t,J=6).

Example 18

20 N-(3,7,11,15-Tetramethyl-2,6,10,14-hexadeca-
tetraenyl)-amyl alcoholamine

The procedure of Example 1 was repeated except that 6.1 g of 3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenoic acid and 3.1 g of amyl alcoholamine were used as starting materials. 7.4 g (yield 95%) of the title compound was obtained as a colorless oil.

Elemental analysis for $C_{25}H_{43}NO_2$

	C	H	N
calculated (%)	77.07	11.13	3.60
30 found (%)	77.01	11.20	3.53

0110397

° Mass (m/z) : 389 (M^+)

° NMR (δ , $CDCl_3$) : 1.4 - 1.8 (6H,m), 1.60 (9H,s),
1.68 (3H,s), 1.9 - 2.2 (12H), 2.13 (3H,d,J=1),
2.44 (1H,s), 3.1 - 3.7 (4H), 5.10 (3H,m),
5.54 (1H,br,s), 5.71 (1H,t,J=5).

Example 19

1-(3,7,11,15-Tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-4-methylpiperazine

The procedure of Example 9 was repeated except that
6.1 g of 3,7,11,15-tetramethyl-2,6,10,14-
hexadecatetraenoic acid and 3.3 ml of 1-methylpiperazine
were used as starting materials. 6.9 g (yield 90%) of
the title compound was obtained as a colorless oil.

Elemental analysis for $C_{25}H_{42}N_2O$

	C	H	N
calculated (%)	77.66	10.95	7.25
found (%)	77.45	11.10	7.30

° Mass (m/z) : 386 (M^+)

° NMR (δ , $CDCl_3$) : 1.60 (9H,s), 1.68 (3H,s),
1.86 (3H,d,J=1), 1.9 - 2.2 (12H),
2.29 (3H,s), 2.2 - 2.45 (4H), 3.4 - 3.7 (4H),
5.10 (3H,m), 5.74 (1H,br,s).

Example 20

1-(3,7,11,15-Tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-4-methyl-hexahydro-1,4-diazepine

The procedure of Example 9 was repeated except that
6.1 g of 3,7,11,15-tetramethyl-2,6,10,14-
hexadecatetraenoic acid and 3.4 g of 4-methyl-1H-
hexahydro-1,4-diazepine were used as starting materials.

0110397

7.5 g (yield 94%) of the title compound was obtained as a colorless oil.

Elemental analysis for $C_{26}H_{44}N_2O$

5		C	H	N
	calculated (%)	77.94	11.07	6.99
	found (%)	77.85	11.10	7.03

° Mass (m/z) : 400 (M^+)

10 ° NMR (δ , $CDCl_3$) : 1.60 (9H,s), 1.68 (3H,s),
1.91 (3H,d,J=1), 1.7 - 2.3 (14H),
2.35 (3H,s), 2.45 - 2.7 (4H), 3.4 - 3.75 (4H),
5.08 (3H,m), 5.78 (1H,br,s).

15 Example 21

N-(3,7,11,15-Tetramethyl-2,6,10,14-hexadeca-
tetraenoyl)-ethanethiolamine

The procedure of Example 1 was repeated except that
6.1 g of 3,7,11,15-tetramethyl-2,6,10,14-
20 hexadecatetraenoic acid and 2.3 g of ethanethiolamine
were used as starting materials. 6.0 g (yield 83%) of
the title compound was obtained as a colorless oil.
Elemental analysis for $C_{22}H_{37}NOS$

25		C	H	N	S
	calculated (%)	72.67	10.26	3.85	8.82
	found (%)	72.55	10.31	3.80	8.91

° Mass (m/z) : 363 (M^+)

30 ° NMR (δ , $CDCl_3$) : 1.60 (9H,s), 1.69 (3H,s),
1.9 - 2.3 (13H), 2.15 (3H,s), 2.5 - 2.8 (2H),
3.3 - 3.6 (2H), 5.11 (3H,m), 5.57 (1H,br,s),
5.85 (1H,br).

0110397

Example 223-(3,7,11,15-Tetramethyl-2,6,10,14-hexadeca-
tetraenoylamino)-1-ethylpiperidine

The procedure of Example 9 was repeated except that
 5 6.1 g of 3,7,11,15-tetramethyl-2,6,10,14-
 hexadecatetraenoic acid and 3.8 g of 3-amino-1-
 ethylpiperidine were used as starting materials. 7.4 g
 (yield 90%) of the title compound was obtained as a pale
 yellow oil.

10 : Elemental analysis for $C_{27}H_{46}N_2O$

	C	H	N
calculated (%)	78.20	11.18	6.76
found (%)	78.41	11.21	6.70

15

° Mass (m/z) : 414 (M^+)

° NMR (δ , $CDCl_3$) : 1.04 (3H,t), 1.58 (12H,s),
 1.66 (3H,s), 1.7 - 2.5 (22'H,m), 4.08 (1H,m),
 5.08 (3H,m), 5.56 (1H,s), 6.04 (1H,br).

20

Example 232-(3,7,11,15-Tetramethyl-2,6,10,14-hexadeca-
tetraenoylaminoethyl)-1-ethylpyrrolidine

The procedure of Example 9 was repeated except that
 25 6.1 g of 3,7,11,15-tetramethyl-2,6,10,14-
 hexadecatetraenoic acid and 3.8 g of 2-aminomethyl-1-
 ethylpyrrolidine were used as starting materials. 7.8 g
 (yield 94%) of the title compound was obtained as a
 brown oil.

30

0110397

Elemental analysis for $C_{27}H_{46}N_2O$

	C	H	N
calculated (%)	78.20	11.18	6.76
5 found (%)	78.39	11.20	6.74

° Mass (m/z) : 414 (M^+)° NMR (δ , $CDCl_3$) : 1.40 (3H,t), 1.58 (9H,s),

1.64 (3'H,s), 1.8 - 2.2 (20H,m),

10 2.6 - 3.4 (3H,m), 3.60 (3H,m), 5.08 (3H,m),

5.70 (1H,s), 7.76 (1H,br).

15

20


25

30

0110397

Example 24

2-(3',7',11',15'-Tetramethyl-2',6',10',14'-
hexadecatetraenoylamino)-1-ethylpiperidine



3.1 ml of triethylamine was added to a
10 solution of 6.1 g of 3,7,11,15-tetramethyl-2,6,10,14-
hexadecatetraenoic acid in tetrahydrofuran. 2.1 ml
of ethyl chlorocarbonate was added dropwise thereto

15

20

25

30

0110397

under stirring and under cooling with ice. The mixture was stirred for 15 min.

2.8 g of 2-amino-1-ethylpiperidine was added thereto and the mixture was stirred at room temperature for 30 min and poured in ice-water. It was then extracted with ethyl acetate and washed with water. After drying over magnesium sulfate followed by distillation of the solvent, the resulting reaction mixture was treated by silica gel column chromatography to obtain 7.3 g (yield: 86%) of the title compound as a colorless oil.


Elemental analysis for $C_{27}H_{46}ON_2$

	C	H	N
calculated (%)	78.20	11.18	6.76
found (%)	78.41	10.97	6.50

- Mass (m/z): 414 (M^+)
- NMR (δ ; $CDCl_3$): 1.04 (3H, t, J=8), 1.59 (9H, s), 1.62 (3H, s), 1.9 - 2.6 (25H, m), 4.10 (1H, br, d), 5.08 (3H, br, t), 5.56 (1H, br, s), 6.10 (1H, br, d)

Example 25

2-(3',7',11',15'-Tetramethyl-2',6',10',14'-hexadecatetraenoylaminoethyl)-1-ethylpyrrolidine



0110397

3.1 ml of triethylamine was added to a solution of 6.1 g of 3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenoic acid in tetrahydrofuran. 2.1 ml of ethyl chlorocarbonate was added dropwise thereto under stirring and under cooling with ice. The mixture was stirred for 15 min.

2.8 g of 2-aminomethyl-1-ethylpyrrolidine was added thereto and the mixture was stirred at room temperature for 30 min and poured in ice-water. It was then extracted with ethyl acetate and washed with water. After drying over magnesium sulfate followed by concentration, the resulting concentrate was treated by silica gel column chromatography to obtain 7.5 g (91%) of the title compound.

Elemental analysis for $C_{27}H_{46}ON_2$

	C	H	N
20 calculated (%)	78.20	11.18	6.76
found (%)	78.36	11.00	6.75


° Mass (m/z): 414 (M^+)

25 ° NMR (δ , $CDCl_3$): 1.40 (3H, t, J=7), 1.60 (9H, S), 1.64 (3H, S), 1.8 - 2.3 (19H, m), 2.5 - 3.3 (4H, m), 3.4 - 3.8 (3H, m), 5.08 (3H, m), 5.68 (1H, br, S), 7.76 (1H, br)

0110397

Example 26

N-(3,7,11,15-Tetramethyl-2,6,10,14-hexadecatetra-
enoylaminoethyl)-pyrrolidine



3.1 ml of triethylamine was added to a solution of 6.1 g of 3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenoic acid in tetrahydrofuran. 2.1 ml of ethyl chlorocarbonate was added dropwise thereto under stirring and under cooling with ice and the mixture was stirred for 15 min.

2.3 g of 1-aminoethylpyrrolidine was added thereto and they were stirred at room temperature for 30 min and then poured in ice-water. It was then extracted with ethyl acetate and washed with water. After drying over magnesium sulfate followed by concentration, the resulting concentrate was treated by silica gel column chromatography to obtain 7.5 g (yield: 95%) of the title compound.

Elemental analysis for $C_{26}H_{44}ON_2$

	C	H	N
calculated (%)	77.94	11.07	6.99
found (%)	77.72	11.13	6.87

° Mass (m/z): 400 (M^+)


° NMR (δ , $CDCl_3$): 1.60 (9H, s), 1.65 (3H, s),

0110397

1.8 - 2.2 (19H, m), 3.0 - 3.8 (8H, m),
4.76 (3H, m), 5.70 (1H, br, S),
7.50 (1H, br)

Example 27

5 N-(3,7,11,15-Tetramethyl-2,6,10,14-hexadecatetra-
enoylaminoethyl)-morpholine



3.1 ml of triethylamine was added to a solution
of 6.1 g of 3,7,11,15-tetramethyl-2,6,10,14-hexade-
catetraenoic acid in tetrahydrofuran. 2.1 ml of
15 ethyl chlorocarbonate was added dropwise thereto
under stirring and under cooling with ice and the
mixture was stirred for 15 min.

2.6 g of 1-aminoethylmorpholine was added there-
20 to and the mixture was stirred at room temperature for
30 min and then poured in ice-water. It was then ex-
tracted with ethyl acetate and washed with water.
After drying over magnesium sulfate followed by
25 concentration, the resulting concentrate was treated
by silica gel column chromatography to obtain 7.6 g
(91%) of the title compound.

Elemental analysis for $C_{26}H_{44}O_2N_2$

30		C	H	N
	calculated (%)	74.95	10.65	6.72
	found (%)	74.90	10.61	6.52

0110397

° Mass (m/z): 416 (M^+)

° NMR (δ , $CDCl_3$): 1.60 (9H, s), 1.68 (3H, s),


1.8 - 2.1 (15H, m), 2.4 - 2.6 (6H, m),

3.1 - 3.5 (2H, m), 3.6 - 3.8 (4H, m),

5 5.10 (3H, m), 5.54 (1H, br, s), 5.96 (1H, br)

Example 28

10 N-(3,7,11,15-Tetramethyl-2,6,10,14-hexadecatetra-
enoyl)-N',N'-dimethylethylenediamine hydrochloride



15

3.1 ml of triethylamine was added to a solution of 6.1 g of 3,7,11,15-tetramethyl-2,6,10,14-hexadecatetra-
enoic acid in tetrahydrofuran. 2.1 ml of ethyl chloro-
20 carbonate was added thereto under stirring and under cooling with ice and the mixture was stirred for 15 min.

1.7 g of N,N-dimethylethylenediamine was added thereto and the mixture was stirred at room temperature
25 for 30 min and then poured in ice-water. It was then extracted with ethyl acetate and washed with water. After drying over magnesium sulfate followed by concentration, the resulting concentrate was treated by
30 silica gel column chromatography to obtain N-(3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenoyl)-N',N'-dimethylethylenediamine. 20 ml of a 1.5 M solution of hydrogen

0110397

chloride in ethyl acetate was added thereto. After 1 h, the product was concentrated to obtain 7.4 g (90%) of the title compound.

Elemental analysis for $C_{24}H_{43}ON_2Cl$

5		C	H	N	Cl
	calculated (%)	70.12	10.54	6.81	8.62
	found (%)	70.01	10.60	6.95	8.77


° NMR (δ , $CDCl_3$): 1.62 (9H, s), 1.68 (3H, s),
 10 1.8 - 2.2 (16H, m), 2.92 (3H, s),
 2.96 (3H, s), 3.2 - 3.8 (4H, m),
 5.10 (3H, m), 5.76 (1H, br, s),
 7.80 (1H, br, s)

15

Example 29

N-(3,7,11,15-Tetramethyl-2,6,10,14-hexadecatetra-
 enoic-N',N'-diethylethylenediamine

20



3.1 ml of triethylamine was added to a solution
 25 of 6.1 g of 3,7,11,15-tetramethyl-2,6,10,14-hexadecatetra-
 enoic acid in tetrahydrofuran. 2.1 ml of ethyl chloro-
 carbonate was added dropwise thereto under stirring and
 under cooling with ice and the mixture was stirred for
 30 15 min.

0110397

2.3 g of N,N-diethylethylenediamine was added thereto and the mixture was stirred at room temperature for 30 min and then poured in ice-water. It was then extracted with ethyl acetate and washed with water.

5 After drying over mangesium sulfate followed by concentration, the resulting concentrate was treated by silica gel column chromatography to obtain 7.2 g (89%) of the title compound.

10 Elemental analysis for $C_{26}H_{46}ON_2$

	C	H	N
calculated (%)	77.55	11.52	6.96
found (%)	77.43	11.63	7.01

15 ° Mass (m/z): 402 (M^+)

° NMR (δ , $CDCl_3$): 1.02 (6H, t, J=8), 1.61 (9H, S),

1.68 (3H, S), 1.8 - 2.2 (15H, m),


2.3 - 2.7 (6H, m), 3.1 - 3.4 (2H, m),

20 5.10 (3H, m), 5.54 (1H, br, S),

6.04 (1H, br.)

Example 30

25 N'-(3,7,11,15-Tetramethyl-2,6,10,14-hexadecatetra-
enoyl)-2-aminopyridine



0110397

The same procedure as in Example 1 was repeated except that 6.1 g of 3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenoic acid and 1.9 g of 2-aminopyridine were used as the starting materials. 7.0 g (92%) of the title compound was obtained as a white wax.

Elemental analysis for $C_{25}H_{36}ON_2$


	C	H	N
calculated (%)	78.90	9.54	7.36
10 found (%)	78.90	9.44	7.32

° Mass (m/z): 380 (M^+)

° NMR (δ , $CDCl_3$): 1.56 (9H, s), 1.65 (3H, s),
1.8 - 2.3 (15H, m), 5.05 (3H, m),
15 5.75 (1H, br, s), 7.20 (1H, br.),
8.1 - 8.6 (4H, m).

Example 31

20 N-(3,7,11,15-Tetramethyl-2,6,10,14-hexadecatetra-
enoyl)-imidazole



The same procedure as in Example 1 was repeated except that 6.1 g of 3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenoic acid and 1.4 g of imidazole were used as the starting materials. 6.5 g (91%) of the title compound was obtained as a colorless oil.

0110397

Elemental analysis for $C_{23}H_{34}ON_2$


	C	H	N
calculated (%)	77.92	9.67	7.90
found (%)	77.85	9.71	7.92

° Mass (m/z): 354 (M^+)

° NMR (δ , $CDCl_3$): 1.57 (6H, s), 1.62 (3H, s),
 1.66 (3H, s), 1.8 - 2.4 (15H, m),
 5.08 (3H, m), 6.26 (1H, br, s),
 7.04 (1H, br, s), 7.46 (1H, br, s),
 8.12 (1H, s)

Example 32

N-(3,7,11,15-Tetramethyl-2,6,10,14-hexadecatetra-
enoyl)-2-amino-2-ethyl-1,3-propanediol



The same procedure as in Example 1 was repeated
 except that 6.1 g of 3,7,11,15-tetramethyl-2,6,10,14-
 hexadecatetraenoic acid and 2.4 g of 2-amino-1,3-
 propanediol were used as the starting materials.
 7.5 g (92%) of the title compound was obtained as a
 colorless oil.

Elemental analysis for $C_{25}H_{43}O_3N$

	C	H	N
calculated (%)	74.03	10.69	3.45
found (%)	73.94	10.82	3.40

0110397

° Mass (m/z): 405 (M^+)


° NMR (δ , $CDCl_3$): 0.90 (3H, t, J=8), 1.5 - 1.7 (14H, m),
1.8 - 2.2 (15H, m), 3.4 - 4.2 (6H, m),
5.08 (3H, m), 5.58 (1H, br. S), 5.72 (1H, br.)

5

Example 33

N-(3,7,11,15-Tetramethyl-2,6,10,14-hexadecatetra-
enoyl)-3-amino-1,2-propanediol

10



15

The same procedure as in Example 1 was repeated
except that 6.1 g of 3,7,11,15-tetramethyl-2,6,10,14-
hexadecatetraenoic acid and 1.9 g of 3-amino-1,2-
propanediol were used as the starting materials.

20

6.5 g (86%) of the title compound was obtained as a
white wax.

Elemental analysis for $C_{23}H_{39}O_3N$

25

	C	H	N
calculated (%)	73.16	10.41	3.71
found (%)	73.08	10.45	3.65


30

° Mass (m/z): 377 (M^+)

° NMR (δ , $CDCl_3$): 1.60 (9H, S), 1.65 (3H, S),
1.8 - 2.2 (15H, m), 3.1 - 3.9 (7H, m),
5.07 (3H, m), 5.54 (1H, br. S),
5.84 (1H, br)

Example 34

N-(3,7,11,15-Tetramethyl-2,6,10,14-hexadecatetra-
enoyl)-N',N',N'',N''-tetramethyldiethylenetriamine



The same procedure as in Example 1 was repeated except that 6.1 g of 3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenoic acid and 3.2 g of N',N',N'',N''-tetramethyldiethylenetriamine were used as the starting materials. 8.0 g of the title compound was obtained as a colorless oil.

15 Elemental analysis for $C_{28}H_{51}ON_3$

	C	H	N
calculated (%)	75.45	11.53	9.43
found (%)	75.10	11.50	9.38

20 ° Mass (m/z): 445 (M^+)


° NMR (δ , $CDCl_3$): 1.60 (9H, s), 1.66 (3H, s),
 1.8 - 2.2 (15H, m), 2.24 (12H, s),
 2.3 - 2.6 (4H, m), 3.3 - 3.6 (4H, m),
 25 5.08 (3H, m), 5.80 (1H, br.s)

Example 35

N-(3,7,11,15-Tetramethyl-2,6,10,14-hexadecatetra-
enoyl)-N',N',N'',N''-tetramethyldiethylenetriamine

30 dihydrochloride

0110397



5 4.0 g of N-(3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenyl)-N',N'',N''',N''''-tetramethyldiethylenetriamine obtained in Example 34 was treated with 20 ml of a 1.5 M solution of hydrogen chloride in ethyl acetate at 5°C for 30 min. The solvent was distilled off. After drying under reduced pressure, 4.6 g of the title compound was obtained as a brown wax.


Elemental analysis for $C_{28}H_{53}ON_3Cl_2$

15		C	H	N	Cl
	calculated (%)	65.09	9.95	8.13	13.73
	found (%)	64.94	9.90	8.20	13.91

° NMR (δ , $CDCl_3$): 1.60 (9H, s), 1.67 (3H, s),
 20 1.8 - 2.3 (15H, m), 2.92 (6H, s),
 2.95 (6H, s), 3.1 - 4.1 (1H, m),
 5.08 (3H, m), 6.06 (1H, br, s)

Example 36

25 N-(3,7,11,15-Tetramethyl-2,6,10,14-hexadecatetraenyl)-N',N'-dimethylaminomethylcarboxamide



30

0110397

5.0 ml of triethylamine was added to a suspension of 5.0 g of N,N-dimethylglycine hydrochloride in dimethyl sulfoxide. 4.4 ml of ethyl chlorocarbonate was added dropwise thereto at 5°C. The mixture was stirred for 30 min.

11 g of 3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenylamine was added thereto and the mixture was stirred at room temperature for 2 h.

The reaction liquid was poured in ice-water. It was then extracted with chloroform and washed with water. After drying over magnesium sulfate followed by distillation of the solvent, the resulting reaction mixture was treated by silica gel chromatography to obtain 9.6 g (71%) of the title compound as a colorless oil.

Elemental analysis for $C_{24}H_{42}ON_2$

	C	H	N
calculated (%)	76.95	11.30	7.48
found (%)	76.83	11.28	7.47


° Mass (m/z): 374 (M^+)

° NMR (δ , $CDCl_3$): 1.62 (9H, s), 1.70 (6H, s),
1.3 - 2.2 (12H, m), 2.28 (6H, s),
2.96 (2H, s), 3.96 (2H, t, J=6),
5.15 (4H, m), 7.05 (1H, br)

0110397

Example 37

N-(3,7,11,15-Tetramethyl-2,6,10,14-hexadecatetra-
enyl)-2-aminoethylcarboxamide



5.6 g of N-t-butyloxycarbonyl- β -alanine was dissolved in 40 ml of tetrahydrofuran. 4 ml of triethylamine was added thereto. 3.2 ml of ethyl chlorocarbonate was added dropwise thereto and the mixture was stirred for 30 min.

8.0 g of 3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenylamine was added thereto and the mixture was stirred at room temperature for 2 h.

The product was after-treated by an ordinary method. After a treatment by silica gel column chromatography, an N-t-butyloxycarbonyl derivative of the title compound was obtained. The product was dissolved in 100 ml of tetrahydrofuran. 30 ml of 5 N hydrochloric acid solution was added thereto. After treatment at room temperature for 5 h followed by the treatment according to silica gel column chromatography, 7.2 g (50%) of the title compound was obtained as a colorless oil.

Elemental analysis for $C_{23}H_{40}ON_2$

30	C	H	N
calculated (%)	76.61	11.18	7.77
found (%)	76.56	11.23	7.70

0110397

Elementary analysis for $C_{23}H_{33}O_2N$

	C	H	N
calculated (%)	77.70	9.36	3.94
found (%)	77.82	9.45	3.89

5 ° Mass (m/z): 355 (M^+)° NMR ($CDCl_3$, δ): 1.58 (3H, s), 1.60 (3H, s),1.63 (3H, s), 1.83 (3H, d, $J=1$),1.9 - 2.3 (8H), 2.68 (1H, t, $J=4$),

10

3.5 - 3.9 (4H, m), 5.10 (2H, m),


6.24 (1H, s), 6.60 (1H, broad),

7.24 (2H, d, $J=8$), 7.50 (2H, d, $J=8$)Example 40

15

N-[4-(2',6',10'-Trimethyl-1',5',9'-undecatrienyl)-benzoyl]diethanolamine

20



25

5 g of 4-(2',6',10'-trimethyl-1',5',9'-undecatrienyl)-benzoic acid and 3.4 g of diethanolamine were treated in the same manner as in Example 39 to obtain 5.4 g (85%) of the title compound as a colorless oil.

Elemental analysis for $C_{25}H_{37}O_3N$

30

	C	H	N
calculated (%)	75.15	9.33	3.51
found (%)	75.09	9.42	3.49


0110397

° Mass (m/z): 399 (M⁺)

° NMR (CDCl₃, δ): 1.57 (3H, s), 1.60 (3H, s),
1.64 (3H, s), 1.84 (3H, d J=2),
1.9 - 2.3 (8H), 3.4 - 4.0 (10H),
5.10 (2H, m), 6.21 (1H, s),
7.20 (2H, d J=8), 7.41 (2H, d J=8)

Example 41

N-[4-(2',6',10'-Trimethyl-1',5',9'-undecatrienyl)-
benzoyl]-N',N'-dimethylethylenediamine



5 g of 4-(2',6',10'-trimethyl-1',5',9'-undecatrienyl)-
benzoic acid and 2 g of N,N-dimethylethylenediamine were
treated in the same manner as in Example 39 to obtain
5.3 g (86%) of the title compound as a colorless oil.

Elemental analysis for C₂₅H₃₈ON₂

	C	H	N
calculated (%)	78.48	10.01	7.32
found (%)	78.53	10.13	7.36

° Mass (m/z): 382 (M⁺)

° NMR (CDCl₃, δ): 1.58 (3H, s), 1.61 (3H, s),
1.65 (3H, s), 1.85 (3H, d J=2),
1.9 - 2.3 (8H), 2.24 (6H, s),
2.48 (2H, t J=5), 3.50 (2H, dt J=5.5)

0110397

° Mass (m/z): 360 (M^+)


° NMR (δ , $CDCl_3$): 1.60 (9H, s), 1.63 (6H, s),
1.8 - 2.4 (16H, m), 3.00 (2H, t, $J=7$),
3.82 (2H, t, $J=6$), 5.08 (4H, m),
6.78 (1H, br.)

5

Example 38

N-(3,7,11,15-Tetramethyl-2,6,10,14-hexadecatetra-
enyl)-N',N'-dimethylethylenediamine

10



4.4 g of N,N-dimethylethylenediamine was dissolved
in 40 ml of dioxane. 5 ml of pyridine was added to the
15 solution. 17.6 g of 3,7,11,15-tetramethyl-2,6,10,14-
hexadecatetraenyl bromide was added thereto and the
mixture was heated under reflux for 2 h and poured in
ice-water. After extraction with n-hexane followed
20 by washing with water and concentration, the resulting
reaction mixture was treated by alumina column chromato-
graphy to obtain 6 g (33%) of the title compound as a
colorless oil.

25 Elemental analysis for $C_{24}H_{44}N_2$

	C	H	N
calculated (%)	79.93	12.30	7.77
found (%)	79.90	12.31	7.83

30

° Mass (m/z): 360 (M^+)


° NMR (δ , $CDCl_3$): 1.52 (9H, s), 1.60 (3H, s),

0110397

1.63 (3H, S), 1.8 - 2.1 (12H, m),
 2.12 (6H, S), 2.2 - 2.8 (5H, m),
 3.07 (2H, br, d, J=8), 5.00 (4H, m)

Example 39

5 N-[4-(2',6',10'-Trimethyl-1',5',9'-undecatrienyl)-
benzoyl]ethanolamine



5 g of 4-(2',6',10'-trimethyl-1',5',9'-undecatrienyl)-
 benzoic acid was dissolved in 30 ml of benzene. 2.3 g of
 thionyl chloride was added to the solution and the mix-
 15 ture was heated under reflux for 30 min and concentrated
 under reduced pressure.

The concentrate was dissolved in 30 ml of ether..
 2 g of ethanolamine was added to the solution under
 20 cooling with ice and the mixture was stirred for 15 min.
 The reaction liquid was washed with 1 N hydrochloric
 acid, then with aqueous sodium bicarbonate solution
 and finally with water. The liquid was dried over
 25 magnesium sulfate and concentrated. The concentrate was
 treated by alumina column chromatography to obtain 4.5 g
 (79%) of the title compound as a white crystal having
 a melting point of 43 to 44.5°C.


0110397

5.10 (2H, m), 6.24 (1H, s),
 6.75 (1H, t J=5), 7.24 (2H, d J=8),
 7.70 (2H, d J=8)

5 Example 42

N-[4-(2',6',10'-Trimethyl-1',5',9'-undecatrienyl)-
benzoyl]-N',N'-dimethylethylenediamine hydrochloride

10



2 g of N-[4-(2',6',10'-trimethyl-1',5',9'-undecatri-
 enyl)benzoyl]-N',N'-dimethylethylenediamine obtained in
 15 Example 41 was treated with 15 ml of a 1.5 M solution
 of hydrogen chloride in ethyl acetate at 5°C for 30 min.
 The solvent was distilled off. After drying under
 reduced pressure, 2.1 g of the title compound was ob-
 20 tained as a yellow, viscous oil.

Elemental analysis for $C_{25}H_{39}ON_2Cl$


	C	H	N	Cl
calculated (%)	71.66	9.38	6.69	8.46
25 found (%)	71.39	9.49	6.47	8.52

° NMR ($CDCl_3$, δ): 1.54 (3H, s), 1.56 (3H, s), 1.60 (3H, s),
 1.80 (3H, s), 1.9 - 2.3 (8H), 2.85 (3H, s),
 2.92 (3H, s), 3.34 (2H, m), 3.80 (2H, m),
 30 5.10 (2H, m), 6.16 (1H, s), 7.18 (2H, d J=8),
 7.90 (2H, d J=8), 8.60 (1H, br), 11.2 (1H, br)

0110397

Example 43

N-[4-(2',6',10'-Trimethyl-1',5',9'-undecatrienyl)-
benzoyl]-N',N'',N''',N''''-tetramethyldiethylenetriamine



5 g of 4-(2',6',10'-trimethyl-1',5',9'-undecatrienyl)-
 10 benzoic acid and 3 g of N',N'',N''',N''''-tetramethyldiethylene-
 triamine were treated in the same manner as in Example
 39 to obtain 6.6 g (91%) of the title compound as a
 colorless oil.

15 Elemental analysis as $C_{29}H_{47}ON_3$

	C	H	N
calculated (%)	76.77	10.44	9.26
found (%)	76.73	10.49	9.38

20 ° Mass (m/z): 453 (M^+)


° NMR: ($CDCl_3$, δ): 1.58 (3H, s), 1.60 (3H, s),
 1.64 (3H, s), 1.84 (3H, d J=2),
 1.9 - 2.3 (8H), 2.24 (12H, s),
 25 2.44 (4H, m), 3.40 (2H, t J=7),
 3.48 (2H, t J=7), 5.10 (2H, m),
 6.23 (1H, s), 7.21 (2H, d J=8),
 7.45 (2H, d J=8)

30

0110397

Example 44

N-[4-(2',6',10'-Trimethyl-1',5',9'-undecatrienyl)-
benzoyl]-N',N',N'',N''-tetramethyldiethylenetriamine
dihydrochloride



2 g of the compound obtained in Example 43 was
 10 treated in the same manner as in Example 42 to obtain
 2.3 g of the title compound as a brown viscous oil.


Elemental analysis as $C_{29}H_{49}ON_3Cl_2$

	C	H	N	Cl
15 calculated (%)	66.14	9.38	7.98	13.46
found (%)	65.88	9.51	7.76	13.38

° NMR ($CDCl_3$, δ): 1.55 (3H, s), 1.57 (3H, s),
 1.60 (3H, s), 1.83 (3H, s),
 20 1.9 - 2.3 (8H), 2.83 (12H bs),
 3.56 (4H, m), 4.00 (4H, m),
 5.10 (2H, m), 6.20 (1H, s),
 7.16 (2H, d J=8), 7.87 (2H, d J=8),
 25 12.1 (2H, br)

Example 45

N-[4-(2',6',10'-Trimethylundecyl)benzoyl]ethanolamine



0110397

5 g of 4-(2',6',10'-trimethylundecyl)benzoic acid and 2 g of ethanolamine were treated in the same manner as in Example 39 to obtain 5.2 g (92%) of the title compound as a colorless oil.

5 Elemental analysis for $C_{23}H_{39}O_2N$


	C	H	N
calculated (%)	76.40	10.87	3.87
found (%)	76.45	10.91	3.81

10 ° Mass (m/z): 361 (M^+)

° NMR ($CDCl_3$, δ): 0.80 (3H, d J=7), 0.84 (9H, d J=7),
0.9 - 1.9 (15H), 2.35 (1H, dd J=12, 8),
2.65 (1H, dd J=12, 8), 2.76 (1H, bs),
15 3.5 - 3.9 (4H), 6.60 (1H, br),
7.16 (2H, d, J=8), 7.65 (2H, d, J=8)

Example 46

20 N-[4-(2',6',10'-Trimethylundecyl)benzoyl]diethanolamine



25

5 g of 4-(2',6',10'-trimethylundecyl)benzoic acid and 3.4 g of diethanolamine were treated in the same manner as in Example 39 to obtain 6.1 g (95%) of the title compound as a colorless oil.

30

0110397

Elemental analysis for $C_{25}H_{43}O_3N$

	C	H	N
calculated (%)	74.03	10.69	3.45
found (%)	73.91	10.76	3.49

5 ° Mass (m/z): 405 (M^+)


° NMR ($CDCl_3$, δ): 0.83 (3H, d, J=7), 0.87 (9H, d, J=7),
 0.9 - 1.9 (15H), 2.33 (1H, dd, J=13, 8),
 2.56 (1H, dd, J=13, 8), 3.4 - 4.0 (10H),
 7.15 (2H, d, J=8), 7.39 (2H, d, J=8)

10

Example 47

N-[4-(2',6',10'-Trimethylundecyl)benzoyl]-N',N'-
dimethylethylenediamine

15



20

5 g of 4-(2',6',10'-trimethylundecyl)benzoic acid
 and 2 g of N,N-dimethylethylenediamine were treated in
 the same manner as in Example 39 to obtain 5.4 g (88%)
 of the title compound as a colorless oil.

Elemental analysis for $C_{25}H_{44}ON_2$

25

	C	H	N
calculated (%)	77.26	11.41	7.21
found (%)	77.08	11.49	7.14

30


0110397

° Mass (m/z): 388 (M⁺)

° NMR: (CDCl₃, δ): 0.82 (3H, d, J=7), 0.85 (9H, d, J=7),
 0.9 - 1.8 (15H), 2.24 (6H, s),
 2.3 - 2.8 (4H, m), 3.49 (2H, dt, J=5, 5)
 6.75 (1H, t, J=5), 7.16 (2H, d, J=7),
 7.68 (2H, d, J=7)

Example 48

N-[4-(2',6',10'-Trimethylundecyl)benzoyl]-N',N'-
dimethylethylenediamine hydrochloride



2 g of the compound obtained in Example 47 was
 treated in the same manner as in Example 42 to obtain 2.1 g
 of the title compound as a colorless, viscous liquid.

Elemental analysis for C₂₅H₄₅ON₂Cl


	C	H	N	Cl
calculated (%)	70.64	10.67	6.59	8.34
found (%)	70.45	10.79	6.53	8.28

° NMR: (CDCl₃, δ): 0.78 (3H, d, J=7), 0.83 (9H, d, J=7),
 0.9 - 1.8 (15H), 2.2 - 2.8 (2H, m),
 2.88 (3H, s), 2.92 (3H, s),
 3.2 - 3.9 (4H), 7.12 (2H, d, J=7),
 7.88 (2H, d, J=7), 8.58 (1H, t, J=5),
 11.8 (1H, br)

0110397

Example 49

N-[4-(2',6',10'-Trimethylundecyl)benzoyl]-N',N',N'',N''-
tetramethyldiethylenetriamine



5 g of 4-(2',6',10'-trimethylundecyl)benzoic acid
 and 3 g of N',N',N'',N''-tetramethyldiethylenetriamine were
 10 treated in the same manner as in Example 39 to obtain
 6.4 g (89%) of the title compound as a colorless oil.

Elemental analysis for $C_{29}H_{53}ON_3$


	C	H	N
15 calculated (%)	75.76	11.62	9.14
found (%)	75.63	11.68	9.09

° Mass (m/z): 459 (M^+)

° NMR ($CDCl_3$, δ): 0.81 (3H, d, J=7), 0.85 (9H, d, J=7),
 20 0.9 - 1.8 (15H), 2.25 (12H, s),
 2.3 - 2.8 (6H, m), 3.41 (2H, t, J=7),
 3.49 (2H, t, J=7), 7.13 (2H, d, J=7),
 7.44 (2H, d, J=7)

25 Example 50

N-[4-(2',6',10'-Trimethylundecyl)benzoyl]-N',N',N'',N''-
tetramethyldiethylenetriamine dihydrochloride



0110397


2 g of the compound obtained in Example 49 was treated in the same manner as in Example 42 to obtain 2.3 g of the title compound as a brown wax.

Elemental analysis for $C_{29}H_{55}ON_3Cl_2$

5		C	H	N	Cl
	calculated (%)	65.39	10.41	7.89	13.31
	found (%)	65.18	10.50	7.77	13.29
	$^{\circ}$ NMR .. (CDCl ₃ , δ)	: 0.79 (3H, d, J=7), -0.85 (9H, d, J=7),			
10		0.9 - 1.8 (15H), 2.1 - 2.8 (2H),			
		2.84 (12H, bs), 3.58 (4H, m),			
		4.00 (4H, m), 6.72 (2H, bs),			
		7.16 (2H, d, J=7), 7.48 (2H, d, J=7)			

15 Example 51

N-[4-(2',6'-Dimethylheptyl)benzoyl]ethanolamine



5 g of 4-(2',6'-dimethylheptyl)benzoic acid and 3 g of ethanolamine used as starting materials were treated in the same manner as in Example 39 to obtain 5.1 g (85%) of the title compound as white crystals.

Melting point ($^{\circ}$ C): 72.5 to 73.5

Elemental analysis for $C_{18}H_{29}O_2N$

30		C	H	N
	calculated (%)	74.18	10.03	4.81
	found (%)	74.24	10.09	4.77


0110397

° Mass (m/z): 291 (M^+)

° NMR ($CDCl_3$, δ): 0.81 (3H, d, J=7), 0.86 (6H, d, J=7),
0.9 - 1.8 (8H), 2.36 (1H, dd, J=13, 8),
2.66 (1H, dd, J=13, 8), 2.85 (1H, br),
3.4 - 3.9 (4H, m), 6.76 (1H, br),
7.16 (2H, d, J=8), 7.66 (2H, d, J=8)

Example 52

N-[4-(2',6'-Dimethylheptyl)benzoyl]-N',N'',N''',N'''-
tetramethyldiethylenetriamine



5 g of 4-(2',6'-dimethylheptyl)benzoic acid and
4.5 g of N',N'',N''',N'''-tetramethyldiethylenetriamine were
treated in the same manner as in Example 39 to obtain
6.6 g (84%) of the title compound as a colorless oil.

Elemental analysis for $C_{24}H_{43}ON_3$

	C	H	N
calculated (%)	73.98	11.12	10.79
found (%)	73.85	11.18	10.66


° Mass (m/z): 389 (M^+)

° NMR ($CDCl_3$, δ): 0.82 (3H, d, J=7), 0.85 (6H, d, J=7),
0.9 - 1.8 (8H), 2.15 (12H, bs),
2.2 - 2.7 (6H, m), 3.44 (4H, b),
7.09 (2H, d, J=8), 7.25 (2H, d, J=8)

0110397

Example 53

N-[4-(2',6',10',14'-Tetramethylpentadecyl)benzoyl]-
ethanolamine



5 g of 4-(2',6',10',14'-tetramethylpentadecyl)benzoic acid and 2 g of ethanolamine were treated in the same manner as in Example 39 to obtain 5.0 g (90%) of the title compound as a colorless oil.

Elemental analysis for $C_{28}H_{49}O_2N$


	C	H	N
15 calculated (%)	77.90	11.44	3.24
found (%)	77.72	11.56	3.31

° Mass (m/z): 431 (M^+)

° NMR : ($CDCl_3$, δ) : 0.80 (3H, d, J=7), 0.85 (12H, d, J=7),
 20 0.9 - 1.8 (22H), 2.32 (1H, dd J=12, 8),
 2.65 (1H, dd J=12, 8), 3.4 - 3.9 (5H, m),
 6.78 (1H, b), 7.12 (2H, d, J=8),
 7.65 (2H, d, J=8)

25 Example 54

N-[4-(2',6',10',14'-Tetramethylpentadecyl)benzoyl]-
N',N',N'',N''-tetramethyldiethylenetriamine



0110397

5 g of 4-(2',6',10',14'-tetramethylpentadecyl)benzoic acid and 3 g of N',N',N'',N''-tetramethyldiethylenetriamine were treated in the same manner as in Example 39 to obtain 6.2 g (91%) of the title compound as a colorless oil.

5 Elemental analysis for $C_{34}H_{63}ON_3$


	C	H	N
calculated (%)	77.06	11.98	7.93
found (%)	76.92	12.04	7.88

10 ° Mass (m/z): 529 (M^+)

° NMR ($CDCl_3$, δ): 0.80 (3H, d, J=7), 0.84 (12H, d, J=7),
0.9 - 1.8 (22H), 2.18 (12H, bs),
2.2 - 2.7 (6H, m), 3.44 (4H, b),
15 7.10 (2H, d, J=8), 7.25 (2H, d, J=8)

Example 55

20 N-{3-[4'-(4'',8''-Dimethylnonyl)phenyl]butanoyl}-
ethanolamine



25 5 g of 3-[4'-(4'',8''-dimethylnonyl)phenyl]butanoic acid and 2 g of ethanolamine were treated in the same manner as in Example 39 to obtain 4.5 g (80%) of the title compound.

30

0110397

Elemental analysis for $C_{23}H_{39}O_2N$


	C	H	N
calculated (%)	76.40	10.87	3.87
found (%)	76.51	11.03	3.85

5 ° Mass (m/z): 361 (M^+)° NMR... (CDCl₃, δ): 0.84 (9H, d, J=7), 1.30 (3H, d, J=7),

0.9 - 1.8 (12H), 2.1 - 2.6 (5H, m),

3.1 - 3.6 (5H, m), 5.83 (1H, b),

10 7.10 (4H, s)

Example 56N-{3-[4'-(4",8"-Dimethylnonyl)phenyl]butanoyl}-N',N',N",N"-tetramethyldiethylenetriamine

20 5 g of 3-[4'-(4",8"-dimethylnonyl)phenyl]butanoic acid was dissolved in 30 ml of benzene. 2.3 g of thionyl chloride was added to the solution. The mixture was heated under reflux for 30 min and then concentrated

25 under reduced pressure.

The concentrate was dissolved in 30 ml of ether. 3 g of N',N',N",N"-tetramethyldiethylenetriamine was added to the solution under cooling with ice and the

30 mixture was stirred for 15 min.

The reaction liquid was washed with 1 N hydrochloric

0110397

acid and then with water, dried over magnesium sulfate and concentrated. The concentrate was treated by column chromatography to obtain 6.1 g (85%) of the title compound as a colorless oil.

5 Elemental analysis for $C_{24}H_{53}ON_3$

	C	H	N
calculated (%)	75.76	11.62	9.14
found (%)	75.59	11.81	9.05


10 ° Mass (m/z) : 459 (M^+)

° NMR .. ($CDCl_3$, δ): 0.84 (9H, d, J=7), 1.28 (3H, d, J=7),
0.9 - 1.8 (12H), 2.10 (6H, s),
2.12 (6H, s), 2.2 - 2.6 (9H),
15 3.1 - 3.5 (4H, m), 7.08 (4H, s)

Example 57

N-{3-[4'-(4",8"-Dimethylnonyl)phenyl]-2-butenoyl}-
ethanolamine

20



25 5 g of 3-[4'-(4",8"-dimethylnonyl)phenyl]-2-butenoyl
acid and 2 g of ethanolamine were treated in the same
manner as in Example 39 to obtain 4.4 g (78%) of the
title compound as white crystals.

30 Melting point ($^{\circ}C$): 48.0 to 49.5

0110397

Elemental analysis for $C_{23}H_{37}O_2N$


	C	H	N
calculated (%)	76.83	10.37	3.90
found (%)	76.71	10.52	3.86

5 ° Mass (m/z): 359 (M^+)

° NMR ($CDCl_3$, δ): 0.84 (9H, d, J=7), 0.9 - 1.8 (12H),
 2.50 (3H, d, J=1), 2.4 - 2.7 (2H, m),
 3.3 - 3.8 (5H, m), 6.02 (1H, q, J=1),
 6.38 (1H, t, J=5), 7.09 (2H, d, J=9),
 7.30 (2H, d, J=9)

Example 58

N-{3-[4'-(4",8"-Dimethylnonyl)-2-butenoyl]-N',N',N",N"-
tetramethyldiethylenetriamine}



20 5 g of 3-[4'-(4",8"-dimethylnonyl)phenyl]-2-butenic
 acid was dissolved in 30 ml of benzene. 5 g of 3-
 [4'-(4",8"-dimethylnonyl)phenyl]-2-butenic acid and 3 g
 of N',N',N",N"-tetramethyldiethylenetriamine were treated
 25 in the same manner as in Example 39 to obtain 5.4 g (75%)
 of the title compound as a colorless oil.

Elemental analysis for $C_{29}H_{51}ON_3$

	C	H	N
30 calculated (%)	76.09	11.23	9.18
found (%)	75.88	11.39	9.15

0110397


° Mass (m/z): 457 (M⁺)

° NMR (CDCl₃, δ): 0.85 (9H, d, J=6), 0.9 - 1.8 (12H),
2.22 (6H, s), 2.28 (9H, s),
2.3 - 2.7 (6H), 3.3 - 3.6 (4H, m),
6.30 (1H, q, J=1), 7.13 (2H, d, J=8),
7.34 (2H, d, J=8)

5

Example 59

N-[2-Methyl-4-(2',5',10'-trimethylundecyl)benzoyl]-
ethanolamine



3 g of 2-methyl-4-(2',6',10'-trimethylundecyl)benzoic
15 acid and 1.5 g of ethanolamine were treated in the same
manner as in Example 39 to obtain 2.9 g (86%) of the
title compound as white crystals.

Melting point (°C): 48.5 to 49.5

20 Elemental analysis for C₂₄H₄₁O₂N


	C	H	N
calculated (%)	76.75	11.00	3.73
found (%)	76.58	11.19	3.76

25 ° Mass (m/z): 375 (M⁺)

° NMR (CDCl₃, δ): 0.80 (3H, d, J=7), 0.84 (9H, d, J=7),
0.9 - 1.8 (15H), 2.25 (1H, dd, J=12, 8),
2.60 (1H, dd, J=12, 8), 2.40 (3H, s),
2.90 (1H, b), 3.4 - 3.9 (4H, m),
30 6.31 (1H, b), 6.8 - 7.3 (3H, m)

Example 60

N-[2-Methyl-4-(2',6',10'-trimethylundecyl)benzoyl]-
N',N',N'',N''-tetramethyldiethylenetriamine



3 g of 2-methyl-4-(2',6',10'-trimethylundecyl)benzoic
 10 acid and 2 g of N',N',N'',N''-tetramethyldiethylenetriamine
 were treated in the same manner as in Example 39 to obtain
 3.4 g (79%) of the title compound as a colorless oil.

Elemental analysis for $C_{30}H_{55}ON_3$


15		C	H	N
	calculated (%)	76.05	11.70	8.87
	found (%)	75.89	11.90	8.92

° Mass (m/z): 473 (M^+).

20 ° NMR ($CDCl_3$, δ): 0.81 (3H, d, J=8), 0.85 (9H, d, J=7),
 0.9 - 1.8 (15H), 1.96 (6H, s),
 2.24 (3H, s), 2.28 (6H, s), 2.1 - 2.7 (6H, m),
 3.1 - 3.7 (4H, m), 6.8 - 7.1 (3H, m)

25 Example 61

N-[2-(Fluoro-4-(2',6',10'-trinethylundecyl)benzoyl]-
ethanolamine



0110397

3 g of 2-fluoro-4-(2',6',10'-trimethylundecyl)benzoic acid and 1.5 g of ethanolamine were treated in the same manner as in Example 39 to obtain 2.9 g (87%) of the title compound as a colorless oil.

5 Elemental analysis for $C_{23}H_{38}O_2NF$

	C	H	N	F
calculated (%)	72.79	10.09	3.69	5.01
found (%)	72.58	10.15	3.67	5.13


10 ° Mass (m/z): 379 (M^+)

° NMR ($CDCl_3$, δ): 0.78 (3H, d, J=7), 0.84 (9H, d, J=7),
0.9 - 1.8 (15H), 2.46 (1H, dd, J=14, 8),
2.80 (1H, dd, J=14, 8), 3.15 (1H, br),
15 3.3 - 3.8 (4H, m), 6.44 (1H, t, J=5),
6.7 - 7.3 (3H, m)

Example 62

N-[2-Chloro-4-(2',6',10'-trimethylundecyl)benzoyl]-

20 N',N',N'',N''-tetramethyldiethylenetriamine



25 2.5 g of 2-chloro-4-(2',6',10'-trimethylundecyl)benzoic acid and 1.5 g of N',N',N'',N''-tetramethyldiethylenetriamine were treated in the same manner as in Example 39 to obtain 3.2 g (91%) of the title compound as a colorless oil.

30

0110397

Elemental analysis for $C_{29}H_{52}ON_3F$


	C	H	N	F
calculated (%)	72.91	10.97	8.80	3.98
found (%)	72.75	10.91	8.93	3.95

5 ° Mass (m/z): 477 (M^+)

° NMR ($CDCl_3$, δ): 0.83 (3H, d, J=7), 0.85 (9H, d, J=7),
 0.9 - 1.7 (15H), 2.04 (6H, s),
 2.28 (6H, s), 2.1 - 2.7 (6H, m),
 3.16 (4H, t, J=7), 6.8 - 7.2 (3H, m)

Example 63

N-[4-(2',6',10'-Trimethylundecyl)-1-naphthoyl]ethanol-
amine



20 5 g of 4-(2',6',10'-trimethylundecyl)-1-naphthoic
 acid and 2 g of ethanolamine were treated in the same
 manner as in Example 39 to obtain 5.2 g (93%) of the title
 compound as white crystals.

25 Melting point ($^{\circ}C$): 61 to 62Elemental analysis for $C_{27}H_{41}O_2N$

	C	H	N
calculated (%)	78.78	10.04	3.40
found (%)	78.85	10.13	3.51


30 ° Mass (m/z): 411 (M^+)

0110397

° NMR (CDCl₃, δ): 0.80 (12H, d, J=7), 0.9 - 1.9 (16H),
 2.66 (1H, dd, J=14, 8), 3.09 (1H, dd,
 J=14, 8), 3.4 - 3.8 (4H, m),
 6.56 (1H, t, J=5), 7.0 - 8.3 (6H, m)

5 Example 64

N-[4-(2',6',10'-Trimethylundecyl)-1-naphthoyl]-N',N',N'',N''-tetramethyldiethylenetriamine



5 g of 4-(2',6',10'-trimethylundecyl)-1-naphthoic
 acid and 3 g of N',N',N'',N''-tetramethyldiethylenetriamine
 15 were treated in the same manner as in Example 39 to obtain
 6.1 g (88%) of the title compound as a colorless oil.


Elemental analysis for C₃₃H₅₅ON₃

	C	H	N
20 calculated (%)	77.74	10.88	8.24
found (%)	77.58	10.95	8.24

° Mass (m/z): 509 (M⁺)

° NMR (CDCl₃, δ): 0.84 (12H, d, J=7), 0.9 - 1.7 (15H),
 25 1.86 (6H, s), 2.0 - 2.3 (2H, m),
 2.36 (6H, s), 2.5 - 3.3 (8H, m),
 7.2 - 8.1 (6H, m)

0110397

Example 65N-[5-(2',6',10'-Trimethylundecyl)-1-naphthoyl]ethanol-amine


5 g of 5-(2',6',10'-trimethylundecyl)-1-naphthoic
 10 acid and 2 g of ethanolamine were treated in the same
 manner as in Example 39 to obtain 5.2 g (93%) of the
 title compound as a colorless oil.

Elemental analysis for $C_{27}H_{41}O_2N$

15		C	H	N
	calculated (%)	78.78	10.04	3.40
	found (%)	78.71	10.12	3.45

° Mass (m/z): 411 (M^+)

20 ° NMR ($CDCl_3$, δ): 0.85 (12H, d, J=7), 0.9 - 2.0 (16H),
 2.70 (1H, dd J=14, 6), 3.10 (1H, dd J=14, 6),
 3.5 - 3.9 (4H, m), 6.47 (1H, t, J=5),
 7.2 - 8.2 (6H, m)

25 Example 66N-[5-(2',6',10'-Trimethylundecyl)-1-naphthoyl]-N',N',N'',N''-tetramethyldiethylenetriamine

0110397

5 g of 5-(2',6',10'-trimethylundecyl)-1-naphthoic acid and 3 g of N',N',N'',N''-tetramethyldiethylenetriamine were treated in the same manner as in Example 39 to obtain 6.0 g (87%) of the title compound as a colorless oil.

5 Elemental analysis for $C_{33}H_{55}ON_3$


	C	H	N
calculated (%)	77.74	10.88	8.24
found (%)	77.72	10.81	8.29

10 ° Mass (m/z): 509 (M^+)

° NMR (CDC $_3$ S): 0.84 (12H, d, J=7), 0.9 - 1.8 (15H),
1.88 (6H, s), 2.1 - 2.3 (2H, m),
2.28 (6H, s), 2.5 - 3.3 (8H, m),
15 7.2 - 8.1 (6H, m)

Example 67

N-[4-(2',6',10'-Trimethylundecyl)-5,6,7,8-tetrahydro-1-naphthoyl]ethanolamine



5 g of 4-(2',6',10'-trimethylundecyl)-5,6,7,8-tetrahydro-1-naphthoic acid and 2 g of ethanolamine were treated in the same manner as in Example 39 to obtain 5.2 g (92%) of the title compound as white crystals.
Melting point (°C): 47 - 48

30 Elemental analysis for $C_{27}H_{45}O_2N$

0110397


	C	H	N
calculated (%)	78.02	10.91	3.37
found (%)	78.16	10.83	3.38

° Mass (m/z): 415 (M⁺)

- 5 ° NMR (CDCl₃, δ): 0.84 (3H, d, J=7), 0.85 (9H, d, J=7),
 0.9 - 1.9 (19H), 2.23 (1H, dd, J=14, 8),
 2.4 - 3.9 (5H, m), 3.10 (1H, br),
 3.3 - 3.8 (4H, m), 6.30 (1H, t, J=5),
 10 6.85 (1H, d, J=8), 7.04 (1H, d, J=8)

Example 68

N-[4-(2',6',10'-Trimethylundecyl)-5,6,7,8-tetrahydro-
1-naphthoyl]-N',N',N'',N''-tetramethyldiethylenetriamine



- 5 g of 4-(2',6',10'-trimethylundecyl)-5,6,7,8-
 20 tetrahydro-1-naphthoic acid and 3 g of N',N',N'',N''-
 tetramethyldiethylenetriamine were treated in the
 same manner as in Example 39 to obtain 5.7 g (82%) of the
 title compound as a colorless oil.

- 25 Elemental analysis for C₃₃H₅₉ON₃

	C	H	N
calculated (%)	77.13	11.57	8.18
found (%)	77.20	11.54	8.25

- 30 ° Mass (m/z): 513 (M⁺)

0110397

° NMR... (CDCl₃, δ): 0.82 (3H, d, J=7), 0.85 (9H, d, J=7),
0.9 - 1.9 (19H), 2.01 (6H, s),
2.30 (6H, s), 2.1 - 3.3 (14H),
6.88 (2H, s)

5

10

15


20

25

30


CLAIMS:

1. A compound of the formula (I):



wherein A, B, Y and Z are each hydrogen, or the pair (1) A and B and/or the pair (2) Y and Z together represent a direct valence bond between the carbon atoms to which they are attached, thereby forming a double bond therebetween; W is a group of -COR or a group of X; and n is zero or an integer of 1 to 4 when W is the group of -COR; n is an integer of 1 to 3 when W is the group of X; R is selected from the group consisting of:


(1) a group of the formula



0110397


wherein R^1 is hydrogen or lower alkyl and m is an integer of from 1 to 5;

(2) a group of the formula




wherein k and l are the same or different and each is an integer of from 1 to 5;

(3) a group of the formula




15 wherein R^2 is hydrogen, lower alkyl or aryl, preferably alkyl or aryl;

(4) a group of the formula



wherein p is an integer of from 0 to 5 and R^3 and R^4 are each hydrogen or lower alkyl, preferably lower alkyl;

(5) a group of the formula




wherein q is an integer of from 1 to 5, R^5 , R^6 and R^7

0110397


are each hydrogen or lower alkyl, preferably lower alkyl, and X is a halogen;

(6) a group of the formula




wherein r is 2 or 3 and R⁸ is lower alkyl;

10 (7) a group of the formula




wherein s is 2 or 3, R⁹ and R¹⁰ are each lower alkyl and X is a halogen;

(8) a group of the formula




wherein D is a group of the formula $\text{---(CH}_2\text{)}_t\text{OH}$, in which
25 t is an integer of from 0 to 5, a group of the formula




wherein u is an integer of from 0 to 5 and R¹¹ and R¹²

0110397

are each hydrogen or lower alkyl, or a group of the formula




wherein v is an integer of from 0 to 5, R^{13} , R^{14} and R^{15}
 10 are each lower alkyl and X is a halogen;
 (9) a group of the formula



wherein R^{16} is hydrogen or lower alkyl and w is an
 integer of from 1 to 5;


(10) a group of the formula



25

wherein R^{17} is hydrogen or lower alkyl and x is an
 integer of from 0 to 5, preferably from 1 to 5; and


(11) a group of the formula




wherein R^{18} is hydrogen or lower alkyl and y is an
 integer of 1 to 5,

X is selected from the group consisting of:


(1) a group of the formula



wherein R^{22} is a group of the formula




wherein R^{23} is hydrogen or lower alkyl and m is an integer of from 1 to 5;
a group of the formula




wherein k and l are the same or different and each is an integer of from 1 to 5;

a group of the formula



wherein p is an integer of from 0 to 5 and R^{24} and R^{25} are each hydrogen or lower alkyl;


and a group of the formula



wherein q and i are each an integer of 1 to 5 and R^{26} , R^{27} , R^{28} and R^{29} are each a lower alkyl, and R^{21} is hydrogen, a lower alkyl or a halogen atom,


0110397

(2) a group of the formula




5 wherein K and L are both hydrogen or represent a direct valence bond the carbon atoms to which they are attached,


(3) a group of the formula




(4) a group of the formula



20 (5) a group of the formula




(6) a group of the formula



wherein a is zero or an integer of 1 to 5, and R^{30} is a lower alkyl,

0110397

(7) a group of the formula



5

wherein b is zero or an integer of 1 to 5 and R^{31} is a lower alkyl,

(8) a group of the formula



10

wherein c is zero or an integer of 1 to 5,

(9) a group of the formula




15

wherein d is zero or an integer of 1 to 5 and R^{32} and R^{33} are each a lower alkyl,


20

(10) a group of the formula



25


(11) a group of the formula



30


wherein R^{34} is a lower alkyl,

(12) a group of the formula




0110397

(13) a group of the formula




(14) a group of the formula



wherein e and f are each an integer of 1 to 5 and R^{35} ,


10 R^{36} , R^{37} and R^{38} are each hydrogen or a lower alkyl,

(15) a group of the formula



15 wherein g is an integer of 1 to 5 and R^{39} and R^{40} are each hydrogen or a lower alkyl, and

(16) a group of the formula



wherein h is an integer of 1 to 5 and R^{41} and R^{42} are each hydrogen or a lower alkyl,

or a pharmaceutically acceptable salt thereof.

25

30

0110397

2. A compound as claimed in Claim 1 wherein W is the group of -COR.

3. A compound as claimed in Claim 1 wherein W is X.

5 4. A pharmaceutical composition which comprises a therapeutically effective amount of a compound as defined in Claim 1, in association with a pharmaceutically acceptable carrier, diluent or vehicle.

10 5. A pharmaceutical composition having anti-PAF activity which comprises a therapeutically effective amount of a compound as claimed in Claim 1, in association with a pharmaceutically acceptable carrier, diluent or vehicle.

15 6. A pharmaceutical composition having antithrombic activity which comprises a therapeutically effective amount of a compound as claimed in Claim 1, in association with a pharmaceutically acceptable carrier, diluent or vehicle.

20

25

30